

2000] Primary 34L20, 05C05; Secondary 34L40

SCHRÖDINGER OPERATOR ON HOMOGENEOUS METRIC TREES: SPECTRUM IN GAPS

ALEXANDER V. SOBOLEV AND MICHAEL SOLOMYAK

ABSTRACT. The paper studies the spectral properties of the Schrödinger operator $\mathbf{A}_{gV} = \mathbf{A}_0 + gV$ on a homogeneous rooted metric tree, with a decaying real-valued potential V and a coupling constant $g \geq 0$. The spectrum of the free Laplacian $\mathbf{A}_0 = -\Delta$ has a band-gap structure with a single eigenvalue of infinite multiplicity in the middle of each finite gap. The perturbation gV gives rise to extra eigenvalues in the gaps. These eigenvalues are monotone functions of g if the potential V has a fixed sign. Assuming that the latter condition is satisfied and that V is symmetric, i.e. depends on the distance to the root of the tree, we carry out a detailed asymptotic analysis of the counting function of the discrete eigenvalues in the limit $g \rightarrow \infty$. Depending on the sign and decay of V , this asymptotics is either of the Weyl type or is completely determined by the behaviour of V at infinity.

1. INTRODUCTION

Counting the number of eigenvalues of a perturbed operator, appearing in the spectral gaps of the unperturbed one, is a classical problem. It was extensively investigated both in the general operator-theoretic setting [2] and in applications to various specific problems of Mathematical Physics (the Hill operator, [13], [19]; the Dirac operator, [15], [4]; the periodic Schrödinger and magnetic Schrödinger operators, [1], [10]; waveguide-type operators, [9], etc.) In this paper we study a new problem of this type, which only recently attracted the attention of specialists: our unperturbed operator is the Laplacian on a homogeneous rooted metric tree Γ . In general, a metric tree is a tree whose edges are viewed as non-degenerate line segments, rather than pairs of vertices, as in the case of the standard (combinatorial) trees. This difference is reflected in the nature of the corresponding Laplacian. For a combinatorial tree this is the discrete Laplacian, whereas the Laplacian $\mathbf{A}_0 = -\Delta$ on a metric tree is represented by a family of the operators $-d^2/dx^2$ on its edges, complemented by the Kirchhoff matching conditions at the vertices. The Laplacian on the homogeneous metric tree has very specific spectral properties which we describe later on in details. In particular, the spectrum has the band-gap structure, with a single eigenvalue of infinite multiplicity in each finite gap. For some other operators on a homogeneous tree, having similar nature, the band-gap structure of the spectrum was established earlier by R. Carlson [11]. In the present paper we study the properties of the perturbed operator $\mathbf{A}_{gV} = \mathbf{A}_0 + gV$ where V is a decaying

Date: 31 July 2001.

1991 Mathematics Subject Classification. [.

real-valued potential, and $g \geq 0$ is a coupling constant. The potential V is assumed to be symmetric, i.e. dependent only on the distance $|x|$ between $x \in \Gamma$ and the root of Γ . This perturbation may produce extra eigenvalues in the gaps of $\mathbf{A}_0$.

For an “observation point” λ inside a gap we denote by $M(\lambda; \mathbf{A}_{gV})$ the number of the eigenvalues of the operator $\mathbf{A}_{\alpha V}$, crossing λ as α varies from 0 to g . For any two points λ_1, λ_2 , $\lambda_1 < \lambda_2$ lying in the same gap, we denote by $N(\lambda_1, \lambda_2; \mathbf{A}_{gV})$ the number of eigenvalues of this operator on the interval (λ_1, λ_2) ; see Sect. 4 for more precise definitions. We are interested in the limiting behaviour of these quantities as the coupling constant g tends to infinity. Compared with other problems of this type, mentioned in the beginning of the Introduction, this problem has many new important features.

The starting point of our investigation is a direct decomposition of the Sobolev space $H^{1,0}(\Gamma)$ on the homogeneous tree Γ . This decomposition is orthogonal with respect to the inner products $\int_{\Gamma} u' \bar{v}' dx$ and $\int_{\Gamma} V(x) u \bar{v} dx$ for all symmetric weight functions V simultaneously. Therefore, it reduces the Laplacian and any Schrödinger operator $\mathbf{A}_V$ with a symmetric potential V . This decomposition was constructed in the paper [17] and has proved very useful for spectral theory of this class of operators. Later it was re-discovered by R. Carlson [12] in a somewhat different setting.

The parts of the operator $\mathbf{A}_V$ in each component of the said orthogonal decomposition turn out to be unitary equivalent to second order differential operators A_{V_k} , $k = 0, 1, \dots$ of the Sturm-Liouville type in the space $L^2(\mathbb{R}_+)$. The potentials V_k are obtained from the original potential V by “shifting” the variable: $V_k(t) = V(t + k)$, $k = 0, 1, \dots$. The operators A_{V_k} act as $-d^2/dx^2 + V_k$, but in contrast to the standard Sturm-Liouville problem, the description of the operator domain of A_{V_k} involves specific matching conditions at the points $t_n = n$, $n = 1, 2, \dots$ (see Section 2). Each component A_{V_k} , $k \geq 1$, enters $\mathbf{A}_V$ with the multiplicity

$$n_0 = 1, \quad n_k = b^k - b^{k-1}, \quad k \geq 1.$$

Here $b \geq 2$ is the integer-valued parameter (the branching number) which characterizes the homogeneous tree completely, see the definition in Subsect. 2.1.

The above orthogonal decomposition plays a central role in our approach. First of all, it allows us to calculate the spectrum of $\mathbf{A}_0$ explicitly (see Theorem 3.3): it consists of the bands $[(\pi(l-1) + \theta)^2, (\pi l - \theta)^2]$, $\theta = \arccos(2(b^{1/2} + b^{-1/2})^{-1})$, and the eigenvalues $\lambda_l = (\pi l)^2$, $l \in \mathbb{N}$. Besides, for $V = 0$ all the components $A_{V_k} = A_0$ are identical, so that the spectrum is of infinite multiplicity. For the perturbed operator this decomposition leads to the representation

$$(1.1) \quad M(\lambda; \mathbf{A}_{gV}) = \sum_{k \geq 0} n_k M(\lambda; A_{gV_k}).$$

A similar formula also holds for $N(\lambda_1, \lambda_2; \mathbf{A}_{gV})$. The presence of the exponentially growing factors n_k hampers the study of these sums. Remembering that the numbers n_k reflect the geometry of the tree, rather than the properties of the potential V , we also

study the counting functions $\widetilde{M}$ and $\widetilde{N}$ ignoring the exponential multiplicities n_k . Precisely, we introduce

$$(1.2) \quad \widetilde{M}(\lambda; \mathbf{A}_{gV}) = \sum_{k \geq 0} M(\lambda; A_{gV_k}),$$

and the quantity $\widetilde{N}$ defined in a similar way.

Clearly, the study of the four functions $M, N, \widetilde{M}, \widetilde{N}$ reduces to that of the individual counting functions $M(\lambda; A_{gV_k}), N(\lambda_1, \lambda_2; A_{gV_k})$ for the operators A_{V_k} . A similar problem for the classical Hill operator was investigated in [13] and, in a more detailed way, in [19]. The general strategy adopted in [19] applies to the operators A_{gV_k} with only minor changes. However, here a new problem emerges: in order to obtain the asymptotic formulas for the sums (1.1), (1.2) one needs an asymptotics of $M(\lambda; A_{gV_k})$ jointly in two parameters: g and k , with a good control of the remainder estimate. In solving this new problem we see the main technical novelty of the paper.

In the paper we obtain several results of rather different type. In this introduction we do not describe them in detail, but concentrate on their principal features. More extended comments are given in the main text. Also, for the sake of discussion we restrict ourselves to the functions $M(\lambda; \mathbf{A}_V)$ and $\widetilde{M}(\lambda; \mathbf{A}_{gV})$ only.

First of all, as in the case of the “classical” Hill operator problem (see [19]), we observe that the behaviour of $M, \widetilde{M}$ is in general radically different for non-positive and non-negative potentials. More precisely, if $V \leq 0$ decays sufficiently quickly at infinity, then the asymptotics is governed by an appropriate Weyl-type formula, and thus it depends on the values of $V(t)$ at all points $t \in \mathbb{R}$, and contains no information on the spectrum of the unperturbed operator. On the contrary, for a non-negative V the asymptotics is determined by the fall-off of V at infinity, and as a rule, depends heavily on some spectral characteristic of the operator $\mathbf{A}_0$. For instance, the behaviour of $\widetilde{M}(\lambda)$ for $V \geq 0$ is described by an integral of the density of states for A_0 . Similar type of asymptotics is also observed for the potentials $V \leq 0$ whose decay at infinity is slow in some specified sense.

In accordance with this general observation our study of the asymptotics is divided in several parts. We begin in Sect. 4 by specifying the conditions on a non-positive symmetric potential V that guarantee the validity of the Weyl-type asymptotics. Further on, we proceed to the cases when the Weyl formula fails and the asymptotics is determined by the behaviour of V at infinity. Here we investigate two types of potentials: power-like and exponentially decaying. In Sect. 5 we state the results for the functions $\widetilde{M}, \widetilde{N}$. A common feature of the asymptotic formulae in Sect. 5 is that virtually all of them contain the density of states for the operator A_0 . It is also worth pointing out that the power-like potentials induce a power-like growth of $\widetilde{M}$ as $g \rightarrow \infty$, whereas the exponential potentials give rise to a logarithmic growth.

The study of the sum (1.1) is postponed until Sect. 9 as it calls for different techniques and is less complete. For the power-like potentials we are able to establish the

asymptotics only for the quantity $\ln M(\lambda)$. For the exponential potentials we provide more detailed asymptotic information. This is possible due to the “self-similarity” of the exponential function. This property allows us to rewrite the formula (1.1) for the function $M(\lambda, \mathbf{A}_{gV})$ in a form which can be interpreted as a Renewal Equation (see [14], [16]). Then the Renewal Theorem ensures a specific asymptotic behaviour of $M(\lambda)$.

Let us briefly outline the contents of the remaining sections. In Sect. 2 we describe the basic orthogonal decomposition of the space $\mathbf{H}^{1,0}(\Gamma)$ and also the parts of the operator $\mathbf{A}_V$ in its components. In Sect. 3 we calculate the spectrum of the Laplacian on Γ . Here we also carry out a detailed analysis of the density of states for the operator A_0 . This function is involved in the asymptotic formulae for $\widetilde{M}$ in the non-Weyl situation. As was mentioned earlier, the study of the perturbed operator $\mathbf{A}_V$ starts in Sect. 4 where the Weyl’s asymptotics is established. The main results on the non-Weyl asymptotics for the functions $\widetilde{M}$ and $\widetilde{N}$ are collected in Sect. 5. Their proofs are given in Sect. 8, preceded by necessary technical preliminaries in Sect. 6, 7. The last Sect. 9 is devoted to the analysis of the functions M, N .

2. LAPLACIAN ON A HOMOGENEOUS TREE AND ITS DECOMPOSITION

2.1. Homogeneous trees and Laplacians on them. Let Γ be a rooted tree with the root o , the set of vertices $\mathcal{V}(\Gamma)$ and the set of edges $\mathcal{E}(\Gamma)$. We suppose that the length of each edge e is equal to 1. Given two points $y, z \in \Gamma$, we write $y \preceq z$ if y lies on the unique simple path connecting o with z ; let $|z|$ stand for the length of this path. We write $y \prec z$ if $y \preceq z$ and $y \neq z$. The relation $\prec$ defines on Γ a partial ordering. If $y \prec z$, we denote

$$\langle y, z \rangle := \{x \in \Gamma : y \preceq x \preceq z\}.$$

In particular, if $e = \langle v, w \rangle$ is an edge, we call v its initial point and say that e emanates from v and terminates at w .

For any $v \in \mathcal{V}(\Gamma)$ the number $|v|$ is a non-negative integer; we call it *generation* of v and denote $\text{Gen}(v)$. For an edge $e \in \mathcal{E}(\Gamma)$ $\text{Gen}(e)$ is defined as the generation of its initial point.

Let an integer $b > 1$ be given. We suppose that for each vertex $v \neq o$ there are exactly b edges emanating from v . We denote them $e_v^1, \dots, e_v^b$ and write e_v^- for the edge terminating at v . We call b the *branching number* of Γ . We always suppose that only one edge emanates from the root o . Thus, the tree Γ is fully determined by the parameter b , and sometimes we use the notation Γ_b . We call any tree Γ_b , with an arbitrary b , *homogeneous*.

The metric topology and the Lebesgue measure on Γ are introduced in a natural way. The space $\mathbf{L}^2(\Gamma)$ is understood as $\mathbf{L}^2$ with respect to this measure.

A function f on Γ belongs to the Sobolev space $\mathbf{H}^1(\Gamma)$ if and only if it is continuous, $f|_e \in \mathbf{H}^1(e)$ for each edge e , and

$$\|f\|_{\mathbf{H}^1(\Gamma)}^2 := \int_{\Gamma} (|f'|^2 + |f|^2) dx < \infty.$$

As usual, $\mathbf{H}^{1,0}(\Gamma) = \{f \in \mathbf{H}^1(\Gamma) : f(o) = 0\}$.

We define the Dirichlet Laplacian $-\Delta$ on Γ as the self-adjoint operator in $\mathbf{L}^2(\Gamma)$, associated with the quadratic form $\int_{\Gamma} |f'|^2 dx$ considered on the form domain $\mathbf{H}^{1,0}(\Gamma)$. It is easy to describe the operator domain $\text{Dom}(\Delta)$ and the action of Δ . Evidently $f \in \text{Dom}(\Delta) \Rightarrow f|_e \in \mathbf{H}^2(e)$ for each edge e and the Euler-Lagrange equation reduces on e to $\Delta f = f''$. In order to describe the matching conditions at a vertex $v \neq o$, denote by f_- the restriction $f|_{e_v^-}$ and by f_j , $j = 1, \dots, b$ the restrictions $f|_{e_v^j}$. The matching conditions at v are

$$f_-(v) = f_1(v) = \dots = f_b(v); \quad f'_1(v) + \dots + f'_b(v) = f'_-(v)$$

where the derivatives on each edge are taken in the direction consistent with the ordering on Γ . The first matching condition comes from the requirement $f \in \mathbf{H}^1(\Gamma)$ which includes the continuity of f , and the second appears as the natural condition in the sense of Calculus of Variations. At the root o we have the boundary condition $f(o) = 0$. It is easy to check that the conditions listed are also sufficient for $f \in \text{Dom}(\Delta)$.

Along with the Laplacian $-\Delta$ we shall be interested also in the Schrödinger operators with a real, bounded and *symmetric* (that is, depending only on $|x|$) potential V :

$$(2.1) \quad \mathbf{A}_V f := -\Delta f + V(|x|)f, \quad f \in \text{Dom}(\Delta).$$

The operator $\mathbf{A}_V$ is self-adjoint. Its quadratic form is given by

$$\mathbf{a}_V[f] = \int_{\Gamma} (|f'|^2 + V(|x|)|f|^2) dx, \quad f \in \mathbf{H}^{1,0}(\Gamma).$$

2.2. The orthogonal decomposition of $\mathbf{L}^2(\Gamma)$. Our techniques is based upon the orthogonal decomposition of $\mathbf{L}^2(\Gamma)$ into a family of subspaces associated with a class of subtrees of Γ . Given a subtree $T \subset \Gamma$, we say that a function $f \in \mathbf{L}^2(\Gamma)$ belongs to $\mathcal{F}_T$ if and only if

$$f = 0 \quad \text{outside } T$$

and

$$(2.2) \quad f(x) = f(y) \quad \text{if } x, y \in T \text{ and } |x| = |y|.$$

Evidently $\mathcal{F}_T$ is a closed subspace of $\mathbf{L}^2(\Gamma)$. It is easy to describe the operator $\mathcal{P}_T$ of orthoprojection onto $\mathcal{F}_T$. To this end, introduce the function

$$b_T(t) = \#\{x \in T : |x| = t\}.$$

In particular,

$$(2.3) \quad b_{\Gamma}(t) = b^k \quad \text{for } k-1 \leq t < k, \quad k \in \mathbb{N}.$$

It is clear that

$$(\mathcal{P}_T f)(x) = \begin{cases} (b_\Gamma(|x|))^{-1} \sum_{y \in T: |y|=|x|} f(y) & \text{for } x \in T; \\ 0 & \text{for } x \notin T. \end{cases}$$

We shall need the subspaces $\mathcal{F}_T$, associated with the subtrees of two following types. Given a vertex v , let

$$T_v = \{x \in \Gamma : x \succeq v\}.$$

Given an edge $e = \langle v, w \rangle$, let

$$T_e = e \cup T_w.$$

In particular, $T_{e_0} = T_o = \Gamma$. For the sake of brevity, for any $v \neq o$ below we use the notation $\mathcal{F}_v$, $\mathcal{F}_v^j$ for $\mathcal{F}_{T_v}$, $\mathcal{F}_{T_{e_v^j}}$. It is clear that the subspaces $\mathcal{F}_v^1, \dots, \mathcal{F}_v^b$ are mutually orthogonal and their orthogonal sum

$$\widetilde{\mathcal{F}}_v = \mathcal{F}_v^1 \oplus \dots \oplus \mathcal{F}_v^b$$

contains $\mathcal{F}_v$. Denote by $\mathcal{F}'_v = \widetilde{\mathcal{F}}_v \ominus \mathcal{F}_v$ the orthogonal complement. The next theorem is a direct consequence of [17], Theorem 5.1 and Lemma 5.2, where a more general class of trees was considered. Later the result was re-discovered by R. Carlson [12], in a slightly different setting. A new detailed exposition, most convenient for our purposes, was recently given in [18].

Theorem 2.1. *Let $\Gamma = \Gamma_b$ for some $b > 1$.*

- (i) *The subspaces $\mathcal{F}'_v$, $o \neq v \in \mathcal{V}(\Gamma)$ are mutually orthogonal and orthogonal to $\mathcal{F}_\Gamma$. Moreover,*

$$(2.4) \quad \mathbf{L}^2(\Gamma) = \mathcal{F}_\Gamma \oplus \sum_{v \in \mathcal{V}(\Gamma) \setminus \{o\}} \oplus \mathcal{F}'_v.$$

- (ii) *Let $V(t)$ be a real, measurable and bounded function on $\mathbb{R}_+$. Then the decomposition (2.4) reduces the Schrödinger operator (2.1), and in particular the Laplacian $-\Delta = \mathbf{A}_0$.*

2.3. Parts of $\mathbf{A}_V$ in the subspaces $\mathcal{F}_\Gamma$, $\mathcal{F}'_v$. According to Theorem 2.1, the description of the spectrum $\sigma(\mathbf{A}_V)$ reduces to the similar problem for the parts of $\mathbf{A}_V$ in the components of the decomposition (2.4). Consider at first the part of $\mathbf{A}_V$ in the subspace $\mathcal{F}_\Gamma$. It is more convenient (and equivalent) to deal with the quadratic form $\mathbf{a}_V$.

It is natural to identify a function $f \in \mathcal{F}_\Gamma$ with the function φ on $\mathbb{R}_+$, such that $\varphi(t) = f(x)$ for $|x| = t$. The operator $\Pi : f \mapsto \varphi$ acts as an isometry of $\mathcal{F}_\Gamma$ onto the weighted space $\mathbf{L}^2(\mathbb{R}_+, b_\Gamma)$ with the norm given by

$$\|\varphi\|_{\mathbf{L}^2(\mathbb{R}_+, b_\Gamma)}^2 = \int_{\mathbb{R}_+} |\varphi(t)|^2 b_\Gamma(t) dt.$$

Then

$$(2.5) \quad \mathbf{a}_V[f] = \int_{\mathbb{R}_+} (|\varphi'(t)|^2 + V(t)|\varphi(t)|^2) b_\Gamma(t) dt, \quad \varphi = \Pi f.$$

Its domain is the weighted Sobolev space $H^{1,0}(\mathbb{R}_+, b_\Gamma)$ whose norm is defined by the quadratic form (2.5) with $V \equiv 1$. The corresponding operator $\mathbf{A}_V \upharpoonright \mathcal{F}_\Gamma$ turns into an operator acting in $L^2(\mathbb{R}_+, b_\Gamma)$. It is not difficult to describe it explicitly, however it is more natural to pass on to the operators acting in the “usual” $L^2(\mathbb{R}_+)$. To this end we make the substitution

$$(2.6) \quad y(t) = b_\Gamma(t)^{1/2} \varphi(t).$$

Then

$$\|y\|_{L^2(\mathbb{R}_+)}^2 = \|\varphi\|_{L^2(\mathbb{R}_+, b_\Gamma)}^2.$$

Since $b_\Gamma(t)$ is a step function, we also have

$$a_V[y] := \mathbf{a}_V[f] = \int_{\mathbb{R}_+} (|y'(t)|^2 + V(t)|y(t)|^2) dt.$$

However, the domain of a_V does not coincide with $H^{1,0}(\mathbb{R}_+)$, since the function $y(t)$ may have jumps at the points $n \in \mathbb{N}$. More exactly, it follows from (2.3) and (2.6) that $\text{Dom}(a_V)$ consists of functions

$$(2.7) \quad y \in H^1(0, 1) \times H^1(1, 2) \times \dots \times H^1(n-1, n) \times \dots$$

such that

$$(2.8) \quad y(0) = 0; \quad y(n+) = b^{1/2} y(n-), \quad \forall n \in \mathbb{N},$$

and

$$(2.9) \quad \int_{\mathbb{R}_+} (|y'(t)|^2 + |y(t)|^2) dt < \infty.$$

The self-adjoint operator in $L^2(\mathbb{R}_+)$, associated with this quadratic form, on each interval $(n-1, n)$, $n \in \mathbb{N}$ acts as

$$A_V y = -y'' + V(t)y.$$

Its domain $\text{Dom}(A_V)$ consists of all functions

$$y \in H^2(0, 1) \times H^2(1, 2) \times \dots \times H^2(n-1, n) \times \dots$$

satisfying the conditions (2.8) and

$$(2.10) \quad y'(n+) = b^{-1/2} y'(n-), \quad \forall n \in \mathbb{N},$$

and also

$$\int_{\mathbb{R}_+} (|y''(t)|^2 + |y(t)|^2) dt < \infty.$$

(Here and in (2.9) it would be more accurate to write $\sum_{n=1}^{\infty} \int_{n-1}^n$ rather than $\int_{\mathbb{R}_+}$.) So we have proved the following

Lemma 2.2. *The part of the operator $\mathbf{A}_V$ in the subspace $\mathcal{F}_\Gamma$ is unitarily equivalent to the operator A_V in $\mathcal{L}^2(\mathbb{R}_+)$.*

Now we turn to the operators $\mathbf{A}_V \upharpoonright \mathcal{F}'_v$, $v \neq o$. It follows from the symmetry properties of the tree Γ and of the potential $V(|x|)$ that all such operators with the same value of $\text{Gen}(v) = k$ can be identified with each other. In order to reduce them to the operators in $\mathcal{L}^2(\mathbb{R}_+)$, introduce the “shifted” potentials

$$(2.11) \quad V_k(t) = V(t + k), \quad t > 0, \quad k = 0, 1, \dots$$

In particular, $V_0 = V$.

Lemma 2.3. *Let $\Gamma = \Gamma_b$ and $v \in \mathcal{V}(\Gamma)$, $\text{Gen}(v) = k > 0$. Then the operator $\mathbf{A}_V \upharpoonright \mathcal{F}'_v$ is unitarily equivalent to the orthogonal sum of $(b-1)$ copies of the operator A_{V_k} .*

For the formal proof, see [18]. On the qualitative level, the result follows from the fact that the restriction of the operator $\mathbf{A}_V$ to the subspace $\widetilde{\mathcal{F}}_v$ reduces to orthogonal sum of b copies of the operator A_{V_k} . The passage to the subspace $\mathcal{F}'_v$ corresponds to the withdrawal of one of these copies.

2.4. The orthogonal decomposition of the operators $\mathbf{A}_V$. Now we are in position to present the final result of this section. Below $A^{[r]}$ stands for the orthogonal sum of r copies of a self-adjoint operator A .

Theorem 2.4. *Let $\Gamma = \Gamma_b$ and the function V be real, measurable and bounded on $\mathbb{R}_+$. Then the Schrödinger operator (2.1) on Γ is unitarily equivalent to the orthogonal sum of the operators acting in $\mathcal{L}^2(\mathbb{R}_+)$:*

$$\mathbf{A}_V \sim A_V \oplus \sum_{k \in \mathbb{N}} \oplus A_{V_k}^{[b^{k-1}(b-1)]}.$$

In particular, for the Laplacian $-\Delta = \mathbf{A}_0$ we get

$$-\Delta \sim A_0^{[\infty]}.$$

This Theorem is a direct consequence of Theorem 2.1 and Lemmas 2.2, 2.3, if one remembers that the total number of vertices of generation k equals b^{k-1} .

3. SPECTRUM OF THE LAPLACIAN ON Γ_b

3.1. The operator A on the whole line. Along with the operator A_0 in $\mathcal{L}^2(\mathbb{R}_+)$ defined as $A_0 y = -y''$ with the boundary and matching conditions (2.8) and (2.10), consider the similar operator, say A , in $\mathcal{L}^2(\mathbb{R})$:

$$(Ay)(t) = -y''(t), \quad t \notin \mathbb{Z},$$

on the analogous domain supplied with the matching conditions

$$(3.1) \quad y(n+) = b^{1/2}y(n-), \quad y'(n+) = b^{-1/2}y'(n-), \quad n \in \mathbb{Z}.$$

The spectrum of A can be found by means of the standard Floquet procedure. The related quasi-periodic problem is

$$y'' + \mu^2 y = 0, \quad y(1+) = e^{i\xi}y(0+), \quad y'(1+) = e^{i\xi}y'(0+),$$

with the parameter (quasi-momentum) $\xi \in [0, 2\pi)$. Taking into account the matching conditions at the point $n = 1$, we can re-write this as

$$(3.2) \quad \begin{aligned} y''(t) + \mu^2 y(t) &= 0, \quad 0 < t < 1; \\ y(1-) &= b^{-1/2}e^{i\xi}y(0+), \quad y'(1-) = b^{1/2}e^{i\xi}y'(0+). \end{aligned}$$

It is quite straightforward to calculate the eigenvalues of the problem (3.2). Introduce the function

$$(3.3) \quad \varphi(\xi) = \arccos \frac{\cos \xi}{R}, \quad R = \frac{b^{1/2} + b^{-1/2}}{2} > 1.$$

Then the numbers μ_l (square roots of eigenvalues) are given by

$$(3.4) \quad \mu_l(\xi) = \begin{cases} \pi(l-1) + \varphi(\xi), & l \text{ is odd,} \\ \pi l - \varphi(\xi), & l \text{ is even,} \end{cases} \quad l \in \mathbb{N}.$$

The function φ is one-to-one on the interval $[0, \pi]$. Later we shall also need its inverse:

$$(3.5) \quad \psi(\mu) = \arccos(R \cos \mu), \quad \mu \in [\varphi(0), \varphi(\pi)] = [\theta, \pi - \theta].$$

where

$$\theta = \arccos(1/R).$$

It follows easily from (3.3) that

$$(3.6) \quad \begin{cases} \psi(\mu) = 2^{1/2}(R^2 - 1)^{1/4}(\mu - \theta)^{1/2} + O(\mu - \theta), & \mu \rightarrow \theta+, \\ \psi(\mu) = \pi - 2^{1/2}(R^2 - 1)^{1/4}(\mu - \pi + \theta)^{1/2} \\ \quad + O(\mu - \pi + \theta), & \mu \rightarrow \pi - \theta - . \end{cases}$$

Define the segments (“bands”)

$$\mathbf{b}_l = \bigcup_{\xi} \mu_l^2(\xi) = [(\pi(l-1) + \theta)^2, (\pi l - \theta)^2], \quad l \in \mathbb{N}$$

and the intervals (“gaps”)

$$(3.7) \quad \mathbf{l}_0 = (-\infty, \theta^2), \quad \mathbf{l}_l = ((\pi l - \theta)^2, (\pi l + \theta)^2), \quad l \in \mathbb{N}.$$

The gaps are labelled so that $\mathbf{l}_l$ separates the bands $\mathbf{b}_l$ and $\mathbf{b}_{l+1}$. The following statement is a direct consequence of the Floquet theory.

Lemma 3.1. *The spectrum of A coincides with the union of the bands $\mathbf{b}_l$, $l = 1, 2, \dots$. On this set the spectrum is of the Lebesgue type and of multiplicity two.*

We shall need also the spectral decomposition of the operator A . To this end, note that

$$\begin{aligned}\zeta_l(t, \xi) &= c_l(\xi) (\cos(\mu_l(\xi)(1-t)) - b^{1/2} e^{i\xi} \cos(\mu_l(\xi)t)), \\ c_l(\xi) &= \sqrt{2(b+1)^{-1}} |\sin \mu_l(\xi)|^{-1}, \quad 0 < t < 1,\end{aligned}$$

is the normalized in $L^2(0, 1)$ eigenfunction of the equation (3.2) corresponding to the eigenvalue $\mu_l^2(\xi)$. It follows from (3.4), (3.3) that $\zeta_l(t, \xi)$ is smooth in ξ on each band $\mathbf{b}_l$. Let us extend each function $\zeta_l(t, \xi)$ to all $t \in \mathbb{R}$ in the following way. Let $\omega_l(t, \xi)$ be the periodic (in t) extension of the function $e^{-it\xi} \zeta_l(t, \xi)$ from the interval $[0, 1)$ to $\mathbb{R}$. Then we define $\zeta_l(t, \xi)$ on the whole of $\mathbb{R}$ by the equation

$$\zeta_l(t, \xi) = e^{it\xi} \omega_l(t, \xi), \quad t \in \mathbb{R}.$$

Let P_l be the spectral projection of A associated with the band $\mathbf{b}_l$. The map

$$(U_l y)(\xi) = (2\pi)^{-1/2} \int_{\mathbb{R}} \overline{\zeta_l(t, \xi)} y(t) dt$$

defines the unitary operator from $L^2(\mathbb{R})$ onto $L^2(-\pi, \pi)$ which diagonalizes AP_l , namely

$$(U_l AP_l y)(\xi) = \lambda_l(\xi) (U_l P_l y)(\xi), \quad \lambda_l(\xi) = \mu_l^2(\xi).$$

The adjoint operator $U^* : L^2(-\pi, \pi) \rightarrow L^2(\mathbb{R})$ is given by

$$(U_l^* z)(t) = (2\pi)^{-1/2} \int_{-\pi}^{\pi} \zeta_l(t, \xi) z(\xi) d\xi = (2\pi)^{-1/2} \int_{-\pi}^{\pi} e^{it\xi} \omega_l(t, \xi) z(\xi) d\xi.$$

Denoting by $[m]$ the operator of multiplication by a scalar function m , we get the spectral decomposition of A in the form

$$(3.8) \quad A = \sum_{l \in \mathbb{N}} U_l^* [\lambda_l] U_l P_l.$$

3.2. Spectrum of the operators A_0 and A_0 .

Theorem 3.2. *The spectrum of the operator A_0 consists of the bands $\mathbf{b}_l$, $l \in \mathbb{N}$ and of the simple eigenvalues $\lambda_l = (\pi l)^2$, $l \in \mathbb{N}$. The corresponding eigenfunctions (normalized in $L^2(\mathbb{R}_+)$) are*

$$(3.9) \quad \begin{aligned} y_l(t) &= c(b) b^{-n/2} \sin(\pi l t), \quad t \in (n-1, n), \quad n \in \mathbb{N} \\ c(b) &= (2(b-1))^{-1/2}. \end{aligned}$$

Proof. The operator A_0 is non-negative, so its spectrum lies on $[0, \infty)$.

1. BANDS. Let D be the operator in $L^2(\mathbb{R}_+)$, defined as follows: its operator domain coincides with the quadratic domain of A_0 , i.e. is defined by (2.7) – (2.9), and for y from this domain

$$(3.10) \quad (Dy)(t) = -iy'(t), \quad t \notin \mathbb{N}.$$

The operator D is closed and its adjoint D^* acts by the same formula (3.10) on the domain consisting of those functions y from the direct product (2.7) which satisfy (2.9) and the matching conditions similar to the ones in (2.8) but with the factor $b^{1/2}$ replaced by $b^{-1/2}$; there is no boundary condition at $t = 0$.

It is easy to see that $A_0 = D^*D$. Along with A_0 , consider the operator DD^* . It acts as $(DD^*y)(t) = -y''(t)$, $t \notin \mathbb{N}$, and its domain is described by the boundary condition $y'(0) = 0$ and the matching conditions

$$y(n+) = b^{-1/2}y(n-), \quad y'(n+) = b^{1/2}y'(n-), \quad n \in \mathbb{N}.$$

According to the general operator theory, the non-zero spectra of the operators $A_0 = D^*D$ and DD^* coincide.

Now, in the definition of the operator A let us replace the matching condition at $t = 0$ by the boundary conditions

$$y(0+) = 0, \quad y'(0-) = 0.$$

The new operator, say A' , splits into the orthogonal sum, $A' = A_0 \oplus A'_0$ where the operator A'_0 acts in $L^2(\mathbb{R}_-)$. Its description is clear from the construction and it is easy to see that the substitution $t \mapsto -t$ reduces A'_0 to DD^* . The essential spectrum of A' is the same as that of A , i.e. $\cup_{l \in \mathbb{N}} \mathfrak{b}_l$. It also coincides with the union of the essential spectra of the operators A_0 and A'_0 , i.e. with each of them. It follows that the essential spectrum of A_0 coincides with the spectrum of A given by Lemma 3.1.

2. EIGENVALUES. The fact that each function $y_l(t)$, cf. (3.9), is the eigenfunction corresponding to the eigenvalue $(\pi l)^2$, can be verified by the direct inspection. Any two solutions satisfying the boundary condition $y(0) = 0$ are proportional to each other, so that this eigenvalues are simple. The direct inspection shows also that $\lambda = 0$ is not an eigenvalue. So it remains to show that any number $\lambda = k^2 > 0$ with $\pi^{-1}k \notin \mathbb{N}$ can not be an eigenvalue. For this purpose we use the explicit formulae for the solutions of the equation

$$(3.11) \quad y''(t) + k^2 y(t) = 0, \quad t \notin \mathbb{N}$$

under the matching conditions (3.1). Namely, let q_1, q_2 be found from the quadratic equation

$$(3.12) \quad q^2 - 2Rq \cos k + 1 = 0$$

where R is defined in (3.3). Suppose that $q_1 \neq q_2$, that is $R|\cos k| \neq 1$. The functions

$$(3.13) \quad \begin{aligned} y_j(t) &= (b^{1/2} \sin k(n-t) + q_j \sin k(t-n+1))q_j^{n-1}, \\ n-1 &< t < n, \quad n \in \mathbb{N}, \quad j = 1, 2 \end{aligned}$$

are solutions of the problem (3.11) – (3.1). Their Wronskian is equal to $y_1 y_2' - y_1' y_2 = b^{1/2}(q_2 - q_1)k \sin k$, so that the solutions y_1, y_2 are linearly dependent only if $\pi^{-1}k \in \mathbb{N}$ which is the excluded case. Any solution satisfying the condition $y(0+) = 0$ is proportional to the function

$$(3.14) \quad \begin{aligned} y_0(t) &= \frac{y_2(t) - y_1(t)}{q_2 - q_1} \\ &= b^{1/2} \frac{q_2^{n-1} - q_1^{n-1}}{q_2 - q_1} \sin k(n-t) + \frac{q_2^n - q_1^n}{q_2 - q_1} \sin k(t-n+1), \\ n-1 &< t < n, \quad n \in \mathbb{N}. \end{aligned}$$

For $\pi^{-1}k \notin \mathbb{N}$ this function does not lie in $\mathbf{L}^2(\mathbb{R}_+)$ and hence, is not an eigenfunction. If $R|\cos k| = 1$, then $q_1 = q_2 = \pm 1$ and it is easy to see that there also are no $\mathbf{L}^2$ -solutions of the problem (3.11) – (3.1), and we are done. $\square$

The result for the operator $\mathbf{A}_0$, that is for the Laplacian on the tree, immediately follows from Theorem 2.4 and Theorem 3.2.

Theorem 3.3. *The spectrum of the operator $\mathbf{A}_0$ is of infinite multiplicity and consists of the bands $\mathbf{b}_l$ and the eigenvalues $\lambda_l = (\pi l)^2$, $l \in \mathbb{N}$.*

We see that the gap $\mathbf{l}_0$ of the operator A is also the gap for $\mathbf{A}_0$, and each gap $\mathbf{l}_l$ of A with $l \geq 1$ splits into two gaps when we turn to the operator $\mathbf{A}_0$:

$$\mathbf{l}_{l,-} = ((\pi l - \theta)^2, (\pi l)^2); \quad \mathbf{l}_{l,+} = ((\pi l)^2, (\pi l + \theta)^2).$$

3.3. Global quasi-momentum and density of states. Define the density of states for the operators A and A_0 as the limit

$$(3.15) \quad \rho(\lambda) = \lim_{|\Delta| \rightarrow \infty} \frac{N_P(\lambda; \Delta)}{|\Delta|},$$

Here we denote $\Delta = (0, L)$, $L \in \mathbb{N}$, and $N_P(\lambda) = \#\{j : \mu_l^2 < \lambda\}$ is the counting function for the operator $By = -y''$ which at the points $1, \dots, L-1$ has the same matching conditions as in (3.1), and also satisfies the boundary conditions

$$y(0) = b^{1/2}y(L), \quad y'(0) = b^{-1/2}y'(L).$$

The subscript P in the notation for the counting function indicates that the operator B has the boundary conditions of this type. If the limit (3.15) exists for these conditions, then it will also exist for any other conditions, and its value will not depend on them. Later, in order to calculate the density of states we shall use the same formula (3.15), but with the counting function of the Dirichlet problem. In this case we do not use any subscripts and simply write $N(\lambda)$.

Let us find eigenvalues of B . Denote $k = \sqrt{\lambda}$, then choose solutions on every interval $(n, n+1)$ in the form

$$y(t) = \alpha_n \cos k(t-n) + \beta_n \sin k(t-n).$$

In view of the matching conditions, we come, with the notations $c = \cos k, s = \sin k$, to the equalities

$$(3.16) \quad \alpha_n = b^{1/2}(\alpha_{n-1}c + \beta_{n-1}s), \quad \beta_n = b^{-1/2}(-\alpha_{n-1}s + \beta_{n-1}c)$$

for $n = 0, \dots, L$. Here we have identified the points with $n = 0$ and $n = L$, so that $\alpha_0 = \alpha_L$ and $\beta_0 = \beta_L$. To solve this system introduce the functions

$$\mathcal{A}(z) = \sum_{n=0}^{L-1} \alpha_n z^n, \quad \mathcal{B}(z) = \sum_{n=0}^{L-1} \beta_n z^n,$$

where z runs over the set of all complex numbers such that $z^L = 1$. Then by (3.16)

$$\mathcal{A}(z) = b^{1/2}z(c\mathcal{A}(z) + s\mathcal{B}(z)), \quad \mathcal{B}(z) = b^{-1/2}z(-s\mathcal{A}(z) + c\mathcal{B}(z)).$$

This system of two equations has non-trivial solution iff its determinant is identically zero:

$$\begin{aligned} & \det \begin{pmatrix} cb^{1/2}z - 1 & sb^{1/2}z \\ -sb^{-1/2}z & cb^{-1/2}z - 1 \end{pmatrix} \\ &= c^2z^2 - cb^{-1/2}z - cb^{1/2}z + 1 + s^2z^2 = z^2 - 2cRz + 1 = 0, \end{aligned}$$

whence

$$R \cos k = Rc = \frac{z + z^{-1}}{2} = \cos \frac{2\pi n}{L}, \quad n = 0, 1, \dots, L-1.$$

It is convenient to write the formulae for the eigenvalues in terms of the function φ defined by (3.3), and the formulae for $N_P(\lambda)$ – in terms of the “global quasi-momentum” $\omega(\lambda)$ which we now define. Namely, $\omega(\lambda) = \pi l$ if $\lambda \in \mathfrak{l}_l$, and for $\lambda \in \mathfrak{b}_l$

$$(3.17) \quad \omega(\lambda) = \begin{cases} \pi(l-1) + \psi(\sqrt{\lambda} - \pi(l-1)), & l \text{ is odd,} \\ \pi l - \psi(\pi l - \sqrt{\lambda}), & l \text{ is even.} \end{cases}$$

Here ψ is the function inverse to φ , cf. (3.5). Evidently

$$(3.18) \quad c\sqrt{\lambda - \theta^2} \leq \omega(\lambda) \leq C\sqrt{\lambda}, \quad \lambda \geq 0.$$

The eigenvalues of the operator B are given by the formulae

$$\begin{aligned} \mu_l(j) &= \pi(l-1) + \varphi\left(\frac{2\pi j}{L}\right), \quad l \text{ odd}; \quad \mu_l(j) = \pi l - \varphi\left(\frac{2\pi j}{L}\right), \quad l \text{ even,} \\ l &\in \mathbb{N}, \quad j = 0, 1, \dots, L-1. \end{aligned}$$

The number $N_P(\lambda)$ depends on the location of λ . For instance, if $\lambda \in \mathfrak{l}_l$, then $N(\lambda) = lL$, so that

$$\rho(\lambda) = \frac{N_P(\lambda, \Delta)}{|\Delta|} = l = \frac{1}{\pi} \omega(\lambda), \quad \lambda \in \mathfrak{l}_l.$$

To cover the case $\lambda \in \mathfrak{b}_l$ we shall consider two options: l is odd or l is even. Suppose first that l is odd and denote $\sqrt{\lambda} = \pi(l-1) + \chi$. Then

$$\begin{aligned} \frac{N_P(\lambda; \Delta)}{L} &= l - 1 + \ell_1(\chi), \\ \ell_1(\chi) &= \frac{1}{L} \# \left\{ j \in [0, L-1) : \varphi\left(\frac{2\pi j}{L}\right) < \chi \right\}. \end{aligned}$$

The term $\ell_1(\chi)$ can be estimated as follows:

$$\left| \ell_1(\chi) - \frac{1}{\pi} \psi(\chi) \right| \leq 2L^{-1}.$$

Hence by (3.17)

$$(3.19) \quad \left| \frac{N_P(\lambda; \Delta)}{L} - \frac{1}{\pi} \omega(\lambda) \right| \leq 2L^{-1}, \quad \forall \lambda > 0.$$

Suppose now that l is even. Denote $\sqrt{\lambda} = \pi l - \chi$. Then

$$\begin{aligned} \frac{N_P(\lambda; \Delta)}{L} &= l - \ell_2(\chi), \\ \ell_2(\chi) &= \frac{1}{L} \# \left\{ j \in [0, L-1) : \varphi\left(\frac{2\pi j}{L}\right) > \chi \right\}. \end{aligned}$$

The term $\ell_2(\chi)$ can be estimated as follows:

$$\left| \ell_2(\chi) + \frac{1}{\pi} \psi(\chi) - 1 \right| \leq 2L^{-1}.$$

Using (3.17) again, we get (3.19). All this results in the formula

$$(3.20) \quad \rho(\lambda) = \frac{1}{\pi} \omega(\lambda)$$

which is well known for the classical Hill operator.

Relying upon the estimate (3.19) we shall prove a similar estimate for the counting function of the Dirichlet problem on an arbitrary interval (R_1, R_2) , not necessarily with integer R_1, R_2 .

Theorem 3.4. *Let $\Delta = (R_1, R_2)$, $R_1, R_2 \in \mathbb{R}$, $R_1 < R_2$. Then the inequality holds:*

$$(3.21) \quad \left| \frac{N(\lambda; \Delta)}{|\Delta|} - \rho(\lambda) \right| \leq C \frac{1 + \sqrt{\lambda}}{|\Delta|}$$

for all $\lambda > 0$, with a universal constant C .

Proof. It is well known that for the Dirichlet realization of the operator $-y''$ on Δ (with no matching conditions inside!) the counting function is controlled by $C|\Delta|\sqrt{\lambda}$, with a universal constant. Since the number of integer points inside Δ is not greater than $|\Delta| + 1$, it follows from the decoupling principle that

$$(3.22) \quad N(\lambda; \Delta) \leq C(|\Delta|\sqrt{\lambda} + |\Delta| + 1).$$

If the length of Δ is small, say $|\Delta| \leq 2$, then (3.21) is implied by (3.18) and (3.22).

Let now $|\Delta| > 2$. Without loss of generality, we may assume $0 \leq R_1 < 1$. Define $L = [R_2]$ (the integer part of R_2), then $L \geq 2$. Let Δ_+ and Δ_- be the intervals $(0, L+1)$ and $(1, L)$ respectively. Then, clearly,

$$N(\lambda; \Delta_-) \leq N(\lambda; \Delta) \leq N(\lambda; \Delta_+),$$

by variation argument. Furthermore, by the decoupling principle,

$$|N(\lambda; \Delta_{\pm}) - N_P(\lambda; \Delta_{\pm})| \leq 4,$$

so that

$$\frac{N_P(\lambda; \Delta_-)}{|\Delta|} - \frac{4}{|\Delta|} \leq \frac{N(\lambda; \Delta)}{|\Delta|} \leq \frac{N_P(\lambda; \Delta_+)}{|\Delta|} + \frac{4}{|\Delta|}.$$

Therefore

$$\left| \frac{N(\lambda; \Delta)}{|\Delta|} - \rho(\lambda) \right| \leq \max_{\pm} \left| \frac{N_P(\lambda; \Delta_{\pm})}{|\Delta|} - \rho(\lambda) \right| + \frac{4}{|\Delta|}.$$

Let us estimate the r.h.s. with the "−" sign. The modulus equals

$$\left| \frac{N_P(\lambda; \Delta_-)}{L-1} \frac{L-1}{|\Delta|} - \rho(\lambda) \right| \leq \left| \frac{N_P(\lambda; \Delta_-)}{L-1} - \rho(\lambda) \right| + \frac{2N_P(\lambda; \Delta_-)}{|\Delta|(L-1)}.$$

In view of (3.19), the first term in the r.h.s. is bounded by $2(L-1)^{-1} \leq 3L^{-1}$ and in view of (3.22), the second term is bounded by

$$\frac{C((L-1)\sqrt{\lambda} + L)}{|\Delta|(L-1)} \leq \frac{2C(\sqrt{\lambda} + 1)}{|\Delta|}.$$

Repeating the same argument for the "+" sign, we arrive at (3.21). $\square$

We conclude this section by discussing the Hölder properties of the global quasimomentum $\omega(\lambda)$ and thus, those of the density of states $\rho(\lambda)$. It is clear from (3.6) that near the edges of the gap $\mathfrak{l} = (\lambda_-, \lambda_+)$ the function ρ has the following behaviour:

$$\rho(\lambda) = \rho(\lambda_{\pm}) \pm \frac{\sqrt{2}}{\pi} \left[\frac{(R^2 - 1)}{\lambda_{\pm}} \right]^{\frac{1}{4}} (\lambda - \lambda_{\pm})^{\frac{1}{2}} + O(\lambda - \lambda_{\pm}), \quad \lambda \rightarrow \lambda_{\pm} + 0 \pm.$$

Together with the formula (3.18) this asymptotics guarantees that

$$(3.23) \quad |\rho(\lambda) - \rho(\lambda_{\pm})| \geq c|\lambda - \lambda_{\pm}|^{\frac{1}{2}}, \quad \lambda \in \mathbb{R},$$

with a constant c depending on l . The formula (3.3) also ensures that the function ψ is $1/2$ -Hölder continuous, i.e.

$$|\psi(\mu_2) - \psi(\mu_1)| \leq C|\mu_2 - \mu_1|^{1/2}, \quad \mu_1, \mu_2 \in [\theta, \pi - \theta].$$

Using (3.18), one can immediately extend this information to the function ω :

$$|\omega(\lambda_2) - \omega(\lambda_1)| \leq C(|\lambda_2^{1/2} - \lambda_1^{1/2}|^{1/2} + |\lambda_2^{1/2} - \lambda_1^{1/2}|), \quad \lambda_1, \lambda_2 \geq \theta,$$

with a constant C independent of λ_1, λ_2 . Later we shall use a less precise, but somewhat more compact consequence of this estimate and (3.20):

$$(3.24) \quad |\rho(\lambda_2) - \rho(\lambda_1)| \leq C|\lambda_2 - \lambda_1|^{1/2}, \quad \lambda_1, \lambda_2 \in \mathbb{R},$$

with a universal constant C .

4. OPERATOR $\mathbf{A}_V$ WITH A DECAYING POTENTIAL. EIGENVALUES IN THE GAPS

4.1. Functions $M(\lambda)$ and $N(\lambda_1, \lambda_2)$. Here we turn to the study of the spectrum of the Schrödinger operators $\mathbf{A}_V$, cf. (2.1), with the real-valued and bounded potential $V(|x|)$ which in an appropriate sense decays as $|x| \rightarrow \infty$. The essential spectrum of $\mathbf{A}_V$ is the same as for the unperturbed operator $\mathbf{A}_0$ (i.e. Laplacian) and therefore, is given by Theorem 3.3. The spectrum of $\mathbf{A}_V$ may include also eigenvalues lying in the gaps of $\mathbf{A}_0$. For their study, the following quantities are standardly used.

Let $\mathcal{C}$ be a self-adjoint operator in a Hilbert space, and let V be its relatively compact perturbation; we denote $\mathcal{C}_V = \mathcal{C} + V$. Suppose that the interval (λ_-, λ_+) is a gap in $\sigma(\mathcal{C})$. Let $\lambda \in (\lambda_-, \lambda_+)$. Define the counting function $M(\lambda; \mathcal{C}_V)$ as the number of eigenvalues of $\mathcal{C}_{\alpha V}$ crossing the point λ while α varies from 0 to 1. In other words,

$$M(\lambda; \mathcal{C}_V) = \sum_{0 < \alpha < 1} \dim \ker(\mathcal{C} + \alpha V - \lambda).$$

If λ coincides with one of the ends of a gap, the function $M(\lambda; \mathcal{C}_V)$ is defined as the corresponding one-sided limit. If V is a perturbation of fixed sign, that is if $V = \pm q$ with a $q \geq 0$, then the function $M(\lambda)$ is increasing (for $V = -q$) or decreasing (for $V = q$) in $\lambda \in (\lambda_-, \lambda_+)$ and increasing in q .

For any subinterval $(\lambda_1, \lambda_2) \subset (\lambda_-, \lambda_+)$ the function $N(\lambda_1, \lambda_2; \mathcal{C}_V)$ is defined as the total multiplicity of eigenvalues of the operator $\mathcal{C}_V$, lying in (λ_1, λ_2) . Note that if $\lambda_- = -\infty$ and $\lambda \leq \lambda_+$, then

$$N(\lambda; \mathcal{C}_V) := N(-\infty, \lambda; \mathcal{C}_V) = M(\lambda; \mathcal{C}_V).$$

According to Theorem 3.3, the following equalities hold:

$$(4.1) \quad M(\lambda; \mathbf{A}_V) = M(\lambda; A_V) + (1 - b^{-1}) \sum_{k \in \mathbb{N}} b^k M(\lambda; A_{V_k}),$$

$$(4.2) \quad N(\lambda_1, \lambda_2; \mathbf{A}_V) = N(\lambda_1, \lambda_2; A_V) + (1 - b^{-1}) \sum_{k \in \mathbb{N}} b^k N(\lambda_1, \lambda_2; A_{V_k}).$$

Recall that the potentials V_k appearing in (4.1), (4.2) were defined in (2.11). These formulae show that the key step to understanding the behaviour of the functions $M(\lambda; \mathbf{A}_V)$, $N(\lambda_1, \lambda_2; \mathbf{A}_V)$ consists in studying the individual terms of the series (4.1), (4.2). More precisely, we need the detailed information about their behaviour depending on the parameter k .

The study of the sums (4.1), (4.2) is hampered by the presence of the exponential factors in their r.h.s. These factors reflect the geometry of the tree rather than the properties of the potential $V(t)$. For this reason, it makes sense to investigate, along with the functions $M(\lambda; \mathbf{A}_V)$, $N(\lambda_1, \lambda_2; \mathbf{A}_V)$, also the functions

$$(4.3) \quad \widetilde{M}(\lambda; \mathbf{A}_V) = \sum_{k \geq 0} M(\lambda; A_{V_k}),$$

$$(4.4) \quad \widetilde{N}(\lambda_1, \lambda_2; \mathbf{A}_V) = \sum_{k \geq 0} N(\lambda_1, \lambda_2; A_{V_k}).$$

For technical reasons, we shall need also the functions M , N for the operators on intervals $\Delta \subseteq \mathbb{R}$. Define $A_{V,\Delta}$ as the operator in $L_2(\Delta)$ acting as $(A_{V,\Delta}y)(t) = -y''(t) + V(t)y(t)$ for $t \notin \mathbb{Z}$, under the zero boundary conditions at each finite end of Δ and the matching conditions (3.1) at the points $n \in \mathbb{Z} \cap \Delta$. In particular, $A_{V,\mathbb{R}_+} = A_V$. Often we use abbreviated notation for the corresponding functions M , N , such as $M(\lambda, V; \Delta)$ or even $M(\lambda; \Delta)$ when the potential V is fixed. Note a convenient relation

$$(4.5) \quad M(\lambda; A_{V_k}, (R_1, R_2)) = M(\lambda; A_V, (R_1 + k, R_2 + k)),$$

which is valid for any $0 \leq R_1 < R_2 \leq \infty$ and integer k 's. This formula is useful when it is more natural to study the dependence of M on the interval Δ than on the potential.

If V is a function of constant sign, then there is a useful relationship between $M(\lambda; A_{V,\Delta})$ and the spectrum of the compact operator

$$(4.6) \quad T(\lambda) = T(\lambda, V, \Delta) = |V|^{1/2}(A_{0,\Delta} - \lambda I)^{-1}|V|^{1/2}, \quad \lambda \notin \sigma(A_{0,\Delta}).$$

Namely, if λ is a regular point of $A_{0,\Delta}$, then

$$(4.7) \quad M(\lambda; V, \Delta) = n_+(1, T(\lambda, V, \Delta)), \quad V \leq 0;$$

$$(4.8) \quad M(\lambda; V, \Delta) = n_-(1, T(\lambda, V, \Delta)), \quad V \geq 0.$$

Here $n_{\pm}(\cdot, T)$ stands for the counting functions of the positive and negative eigenvalues $\pm \lambda_j^{\pm}(T)$ of a compact, self-adjoint operator T , that is

$$n_{\pm}(s, T) = \#\{j : \lambda_j^{\pm}(T) > s\}, \quad s > 0.$$

The equalities (4.7), (4.8) proved very effective in the problems of the type considered, see e.g. [19]. Actually, these are facts of rather general a nature, see e.g. [2], Proposition 1.5.

The following relations have their prototypes in the theory of the perturbed Hill operator, see [19], (2.5) – (2.8). For bounded Δ

$$(4.9) \quad \begin{cases} M(\lambda; V, \Delta) = N(\lambda; V, \Delta) - N(\lambda; 0, \Delta), & V \leq 0, \\ M(\lambda; V, \Delta) = N(\lambda+; 0, \Delta) - N(\lambda+; V, \Delta), & V \geq 0. \end{cases}$$

Further, for any (bounded or unbounded) Δ

$$(4.10) \quad |N(\lambda_1, \lambda_2; V, \Delta) - |M(\lambda_2; V, \Delta) - M(\lambda_1; V, \Delta)|| \leq N(\lambda_1, \lambda_2; 0, \Delta) + 1.$$

One can give a more precise formula: for any two points λ_1, λ_2 such that $N(\lambda_1, \lambda_2; 0, \Delta) = 0$, we obtain from (4.9):

$$(4.11) \quad \begin{cases} N(\lambda_1, \lambda_2; V, \Delta) = M(\lambda_2; V, \Delta) - M(\lambda_1+; V, \Delta), \\ N(\lambda_1, \lambda_2+; V, \Delta) = M(\lambda_1; V, \Delta) - M(\lambda_2; V, \Delta). \end{cases}$$

The next two inequalities are usually referred to as the “decoupling principle”. Let $\Delta_1 = (a, d)$, $\Delta_2 = (d, c)$, $-\infty \leq a < d < c \leq \infty$, and $\Delta = (a, c)$. Then

$$(4.12) \quad |N(\lambda_1, \lambda_2; V, \Delta) - (N(\lambda_1, \lambda_2; V, \Delta_1) + N(\lambda_1, \lambda_2; V, \Delta_2))| \leq 2,$$

$$(4.13) \quad |M(\lambda; V, \Delta) - (M(\lambda; V, \Delta_1) + M(\lambda; V, \Delta_2))| \leq 2.$$

The proofs of the relations (4.9) – (4.13) are either straightforward, or are based upon standard facts from the perturbation theory. Note that the number 1 rather than 2 stands in the r.h.s of the inequalities [19] (2.7) and (2.8) whose analogs are the above inequalities (4.12), (4.13). This difference appears due to the nature of the matching conditions at the points $n \in \mathbb{Z}$. If $d \notin \mathbb{Z}$, one can replace 2 by 1 in (4.12) and (4.13).

4.2. Individual Weyl asymptotics. The material presented in this subsection, is a minor refinement of [19], Theorems 3.2 and 3.3. We give it here for the operators we need in this paper (that is, the functions in the domains of the operators considered are subject to the matching conditions (3.1)). However, it is useful to keep in mind that the results of Theorem 4.2(i) and of Theorem 4.3 hold also for the usual Hill operator.

For a real-valued function V on $\mathbb{R}_+$, introduce the quantity

$$J(V) = \sum_{n \in \mathbb{Z}} \left(\int_{2^{n-1}}^{2^n} t |V(t)| dt \right)^{1/2}.$$

Consider the operator on $L^2(\mathbb{R}_+)$:

$$(4.14) \quad K_V y = -y'' + Vy, \quad y \in H^2(\mathbb{R}_+), \quad y(0) = 0.$$

In contrast to the operator A_V , the description of K_V involves no matching conditions, and the quadratic domain of K_V is $H^{1,0}(\mathbb{R}_+)$.

The following estimate and asymptotics are particular cases of the results of [8], Sect. 6; see also expositions in [5] and [6]. A close result was obtained earlier in [7], Theorems 4.18, 4.19.

Proposition 4.1. *Let $J(V) < \infty$. Then the negative spectrum of the operator K_V is finite and there exists an absolute constant $C > 0$ such that*

$$(4.15) \quad M(0; K_V) \leq C J(V_-).$$

Besides, let $g > 0$ be the large parameter and $V_- \not\equiv 0$. The function $M(0; K_{gV})$ satisfies Weyl's asymptotics

$$(4.16) \quad \lim g^{-1/2} M(0; K_{gV}) = \pi^{-1} \int_{\mathbb{R}_+} \sqrt{V_-(t)} dt, \quad g \rightarrow \infty.$$

If $|V(t)|$ monotonically decreases, then by Hölder's inequality

$$(4.17) \quad J(V)/\sqrt{6} \leq \int_{\mathbb{R}_+} |V(t)|^{1/2} dt \leq \sqrt{6} J(V).$$

Hence, for monotone $|V(t)|$ the function $M(0; K_{gV})$ is controlled by the r.h.s. of its asymptotics given by (4.16).

Along with $J(V)$, introduce the functional

$$\tilde{J}(V) = \left(\int_0^1 |V(t)| dt \right)^{1/2} + \sum_{n=1}^{\infty} \left(\int_{2^{n-1}}^{2^n} t |V(t)| dt \right)^{1/2}.$$

Clearly $J(V) \leq c \tilde{J}(V)$, therefore in the r.h.s of (4.15) $J(V)$ can be replaced by $\tilde{J}(V)$. Compared with Theorem 4.1, its corollary with $\tilde{J}(V)$ in the r.h.s ignores the fact that due to the Dirichlet condition at 0 the potential V need not be integrable at this point. Still, this corollary is quite convenient provided one is dealing with V integrable at 0.

Present also an estimate for $M(-1; A_V)$; we need it in the course of the proof of Theorem 4.2 below.

$$(4.18) \quad M(-1; A_V) \leq C \Theta(V) := C \sum_{n \in \mathbb{N}} \left[\int_{n-1}^n |V(t)| dt \right]^{1/2}.$$

For the proof, one splits $\mathbb{R}_+$ into the union of the intervals $(n-1, n]$ and applies to each interval the well known eigenvalue estimate for the equation $-y'' + y = \lambda V y$ with the Neumann boundary conditions. This is exactly the way in which the same estimate for $M(-1; K_V)$ was proved in [3].

It follows from Hölder's inequality that $\Theta(V) \leq \tilde{J}(V)$. However, the functional $\Theta(V)$ can not be estimated by $J(V)$. Note that similarly to (4.17), for a decreasing $|V|$ we

have

$$(4.19) \quad \tilde{J}(V) \leq \left[\int_0^1 |V(t)| dt \right]^{1/2} + \sqrt{6} \int_{\mathbb{R}_+} \sqrt{|V(t)|} dt.$$

Theorem 4.2. *Let V be a function with a fixed sign and let $\tilde{J}(V) < \infty$.*

- (i) *Suppose that $\lambda \in \bar{l}_l$ where l_l is one of the gaps (see (3.7)). Then, given an interval $\Delta \subseteq \mathbb{R}$, the estimate*

$$(4.20) \quad M(\lambda; V, \Delta) \leq C(\tilde{J}(V) + 1)$$

holds, where the constant $C = C(l)$ does not depend on $\lambda \in \bar{l}_l$ and V .

- (ii) *Suppose in addition that $\Delta = \mathbb{R}_+$ (so that $A_{V,\Delta} = A_V$), and that*

$$(4.21) \quad \lambda \in I \text{ where } I \subset \begin{cases} \bar{l}_0, & l = 0, \\ \bar{l}_l \setminus \{\lambda_l\}, & l \geq 1, \end{cases} \text{ is a closed interval.}$$

Then

$$(4.22) \quad M(\lambda; A_V) \leq C' \tilde{J}(V)$$

with a constant C' uniform in $\lambda \in I$. In particular, for $V \leq 0$ the estimate (4.22) is uniform in $\lambda \leq \theta^2$.

Proof. The proof of (i) follows the scheme suggested in [19]. For this reason, we only outline the necessary changes in the argument. To be definite, we suppose that $V \leq 0$ and that l (the index of the gap) is even. We start with the spectral decomposition (3.8) of the operator $A = A_{0,\mathbb{R}}$ on the whole line. Set

$$\begin{aligned} \lambda(\xi) &= \lambda_{l+1}(\xi), \quad P = P_{l+1}, \\ P^+ &= \sum_{j>l} \oplus P_j, \quad Q = P^+ - P, \quad U = U_{l+1}. \end{aligned}$$

As in [19], Section 4, the estimating of $M(\lambda; V, \Delta)$ is reduced to the problem of eigenvalue estimates for the operators

$$\begin{aligned} T_1(\lambda) &= |V|^{1/2}(A - \lambda I)^{-1}P|V|^{1/2}, \\ T_2(\lambda) &= |V|^{1/2}(A - \lambda I)^{-1}Q|V|^{1/2}. \end{aligned}$$

Since $\|(A + I)(A - \lambda I)^{-1}Q\| \leq C_l$ (actually, $C_l = O(l)$), we have

$$n_+(1, T_2(\lambda)) \leq n_+(C^{-1}, (|V|^{1/2}(A + I)^{-1}|V|^{1/2}) = M(-1; A + C_l V).$$

To the latter quantity the estimate (4.18) applies, and we obtain

$$(4.23) \quad n_+(1, T_2(\lambda)) \leq C' \Theta(V).$$

To the operator $T_1(\lambda)$ the argument of [19] applies without changes. Indeed, the nature of the operator U in our case is the same as in the case of periodicity coming from

a potential. This allows to reduce the problem to estimating the counting function $N(\lambda; K_V)$ for the operator K_V defined in (4.14). Then using the bound (4.15), we arrive at the inequality $n(1, T_1(\lambda)) \leq C(\tilde{J}(V) + 1)$ which, in combination with (4.23) leads to (4.20).

(ii) Again, for definiteness, we prove the result for the non-positive potentials. Let $\lambda = k^2 \in I, k > 0$. In the case $l \geq 1$ assume temporarily that $\lambda \neq (\pi l - \theta)^2$ and $\lambda \neq (\pi l + \theta)^2$, so that $k \in (\pi l - \theta, \pi l) \cup (\pi l, \pi l + \theta)$. In the case $l = 0$ assume that $\lambda \in (0, \theta^2)$, so that $k \in (0, \theta)$. The roots q_1, q_2 of the equation (3.12) are real and distinct, and $q_1 q_2 = 1$. Let us label them so that $|q_1| < 1 < |q_2|$ and denote $|q_1| = e^{-\sigma}$, then $\sigma > 0$. Consider the solutions y_0 and y_1 (cf. (3.14) and (3.13)) of the problem (3.11) – (3.1). Their Wronskian is $W(y_0, y_1) = y_0 y_1' - y_0' y_1 = -b^{1/2} k \sin k \neq 0$, so that y_0, y_1 are linearly independent. On the interval $(n-1, n)$ the function $y_0(t)$ satisfies the inequality

$$\begin{aligned} |y_0(t)| &\leq nb^{1/2}(q_2^{n-2} + q_2^{n-1}) \leq 2nb^{1/2}e^{\sigma t}, \quad n > 1; \\ |y_0(t)| &\leq |\sin kt| \leq kt, \quad n = 1. \end{aligned}$$

For $y_1(t)$ we have

$$|y_1(t)| \leq (b^{1/2} + 1)q_1^{n-1} \leq q_2(b^{1/2} + 1)e^{-\sigma t}, \quad t > 0.$$

Note also that $|q_2| = R|\cos k| + (R^2 \cos^2 k - 1)^{1/2} \leq 2R$. So we see that the inequalities

$$(4.24) \quad |y_0(t)| \leq cte^{\sigma t}, \quad |y_1(t)| \leq ce^{-\sigma t}, \quad t > 0$$

hold uniformly in $\lambda \in I$. The solution y_0 satisfies the boundary condition $y_0(0+) = 0$.

Given a function $f \in L^2(\mathbb{R}_+)$, the solution of the non-homogeneous equation on $\mathbb{R}_+$:

$$y''(t) + k^2 y(t) = -f(t), \quad t \notin \mathbb{N}; \quad y(0+) = 0$$

satisfying the matching conditions (3.1) for $n \in \mathbb{N}$, is given by

$$y(t) = \int_{\mathbb{R}_+} K(t, s) f(s) ds$$

where

$$W(y_0, y_1)K(t, s) = \begin{cases} y_1(t)y_0(s), & s < t, \\ y_1(s)y_0(t), & t < s. \end{cases}$$

It follows from (4.24) that

$$(4.25) \quad |K(t, s)| \leq c^2 e^{-\sigma|t-s|} \min(s, t)(b^{1/2}k|\sin k|)^{-1} \leq C_1 \sqrt{st}(k|\sin k|)^{-1}, \quad C_1 = c^2 b^{-1/2}.$$

The operator (4.6) (for $\lambda = k^2$ and $\Delta = \mathbb{R}_+$) acts as

$$(T(\lambda)f)(t) = |V(t)|^{1/2} \int_{\mathbb{R}_+} K(t, s)|V(s)|^{1/2} f(s) ds.$$

Under the assumption $\tilde{J}(V) < \infty$ this operator belongs to the Hilbert – Schmidt class. Indeed, by virtue of (4.25)

$$\begin{aligned} & (k \sin k)^2 \iint_{\mathbb{R}_+^2} |K(t, s)|^2 |V(t)| |V(s)| dt ds \\ & \leq C_1^2 \iint_{\mathbb{R}_+^2} st |V(t)| |V(s)| dt ds = C_1^2 \left(\int_{\mathbb{R}_+} t |V(t)| dt \right)^2 \leq C_1^2 \tilde{J}(V)^4. \end{aligned}$$

Since $\|T\| \leq \|T\|_{HS}$ and $n_+(1, T) = 0$ if $\|T\| \leq 1$, the last estimate and (4.7) imply that $M(\lambda; A_V) = 0$ if $\tilde{J}(V)^2 \leq C_1^{-1} k |\sin k|$. In its turn, this and the estimate (4.20) yield the inequality (4.22) with $C' = C(1 + (C_1^{-1} k |\sin k|)^{-1/2})$. By continuity, (4.22) extends to the ends of the gap, i.e. to $k = \pi l \pm \theta$ ($l \geq 1$) or $k = \theta$ ($l = 0$). Since the function $M(\lambda; A_V)$ is monotone in λ , the result for $l = 0$ automatically extends to all $\lambda \leq \theta$. $\square$

As in (4.19) one can simplify the estimate (4.20) if one assumes that $|V|$ is decreasing on the interval $\Delta = (R_1, R_2)$:

$$(4.26) \quad M(\lambda; A_V, \Delta) \leq C \left[\left(\int_{R_1}^{R_1+1} |V(t)| dt \right)^{1/2} + \sqrt{6} \int_{\Delta} \sqrt{|V(t)|} dt + 1 \right].$$

Theorem 4.3. *Let the assumptions of Theorem 4.2 be satisfied and $V \leq 0$. Then the asymptotics*

$$(4.27) \quad \lim g^{-1/2} M(\lambda; gV, \Delta) = \pi^{-1} \int_{\Delta} \sqrt{|V(t)|} dt, \quad g \rightarrow \infty$$

(cf. (4.16)) holds uniformly in $\lambda \in \mathfrak{l}_l$.

Proof. Like in [19], the problem reduces to the case of a finite interval Δ . In view of (4.9) we need to study only the term depending on g . Removal of the matching conditions inside the interval shifts the function $N(\lambda; gV, \Delta)$ no more than by $2|\Delta| + 2$ and therefore, does not affect its asymptotic behaviour. As a result, we come to the operator of the Dirichlet problem on a finite interval for which the asymptotics (4.27) is well known. $\square$

4.3. Weyl asymptotics for $M(\lambda; \mathbf{A}_{gV})$ and $\tilde{M}(\lambda; \mathbf{A}_{gV})$. The results of this subsection follow immediately from Theorems 3.2 and 3.3.

Theorem 4.4. *Let $\Gamma = \Gamma_b$ and let $V(t) \leq 0$ be a bounded measurable function on $\mathbb{R}_+$. Let λ satisfy (4.21). Then*

(i) *If*

$$(4.28) \quad \sum_{k \in \mathbb{N}} b^k \tilde{J}(V_k) < \infty,$$

then the Weyl asymptotics holds for the function $M(\lambda; \mathbf{A}_{gV})$ of the operator (2.1):

$$(4.29) \quad \lim g^{-1/2} M(\lambda; \mathbf{A}_{gV}) = \frac{1}{\pi} \int_{\Gamma} \sqrt{|V(|x|)|} dx, \quad g \rightarrow \infty.$$

(ii) If

$$(4.30) \quad \sum_{k \in \mathbb{N}} \tilde{J}(V_k) < \infty,$$

then

$$(4.31) \quad \lim g^{-1/2} \tilde{M}(\lambda; \mathbf{A}_{gV}) = \frac{1}{\pi} \sum_{k=0}^{\infty} \int_k^{\infty} \sqrt{|V(t)|} dt, \quad g \rightarrow \infty.$$

The above asymptotic formulae are uniform in λ on any closed interval I from (4.21).

Proof. For definiteness, we prove (4.29). The proof of (4.31) is the same.

It follows from (4.1) that

$$(4.32) \quad \begin{aligned} & g^{-1/2} M(\lambda; \mathbf{A}_{gV}) \\ &= g^{-1/2} M(\lambda; A_{gV}) + (1 - b^{-1}) \sum_{k \in \mathbb{N}} b^k g^{-1/2} M(\lambda; A_{gV_k}). \end{aligned}$$

By Theorem 4.3, for each $k \geq 0$

$$g^{-1/2} M(\lambda; A_{gV_k}) \rightarrow \pi^{-1} \int_{\mathbb{R}_+} \sqrt{|V_k(t)|} dt, \quad g \rightarrow \infty.$$

The series (4.28) dominates the series (4.32) and it follows from Lebesgue's theorem on the dominated convergence that

$$\pi g^{-1/2} M(\lambda; \mathbf{A}_{gV}) \rightarrow \int_{\mathbb{R}_+} \sqrt{|V(t)|} dt + (1 - b^{-1}) \sum_{k=1}^{\infty} b^k \int_{\mathbb{R}_+} \sqrt{|V_k(t)|} dt, \quad g \rightarrow \infty.$$

This is equivalent to (4.29). Due to Theorem 4.2 the series (4.28) converges uniformly on I from (4.21), and hence the asymptotics (4.29) is uniform in $\lambda \in I$. $\square$

5. POWER-LIKE AND EXPONENTIAL POTENTIALS

5.1. Weyl asymptotics. Here we show how the theorems in the previous section apply to potentials with a specified rate of decay at infinity. To have a clear distinction between the cases of non-positive and non-negative potentials, we slightly change our notation: we denote the potential by $V = \pm q$ or $V = \mathbf{s}q$ with $\mathbf{s} = \pm 1$, always assuming that $q \geq 0$.

We are interested in two types of potentials: power-like and exponential. More precisely, suppose that $q(t) \leq CQ(t)$ where

$$(5.1) \quad Q(t) = (1+t)^{-2\gamma}, \quad \gamma > 0, \quad \text{or} \quad Q(t) = e^{-2\kappa t}, \quad \kappa > 0.$$

Let us first establish the Weyl type asymptotics for $M(\lambda; \mathbf{A}_{-gq})$ and $\tilde{M}(\lambda; \mathbf{A}_{-gq})$.

Theorem 5.1. *Let Condition (4.21) be fulfilled. Suppose that $q(t) \leq CQ(t)$.*

- (i) *If $Q(t) = (1+t)^{-2\gamma}$ with $\gamma > 2$, then the asymptotic formula (4.31) holds for $\widetilde{M}(\lambda; \mathbf{A}_{-gq})$;*
- (ii) *If $Q(t) = e^{-2\kappa t}$, $\kappa > 0$, then the asymptotic formula (4.31) for $\widetilde{M}(\lambda; \mathbf{A}_{-gq})$ holds. If, in addition, $\kappa > \ln b$, then the asymptotics (4.29) for $M(\lambda; \mathbf{A}_{-gq})$ holds as well.*

These results are uniform in $\lambda \in I$ with a closed interval I from (4.21).

The proof of Theorem 5.1 is based on two elementary Lemmas 5.2 and 5.3 describing individual counting functions. These Lemmas will be also useful in the analysis of the non-Weyl behaviour of the function $\widetilde{M}$.

Recall that by q_k , $k \geq 0$ are denoted the “shifted” potentials $q_k(t) = q(t+k)$, $t > 0$. Remembering the relation (4.5) and a comment after it, we sometimes transfer the dependence on k to the interval Δ . This is why some of the estimates below are stated for intervals Δ depending on an additional parameter R , which plays the role of k , but is not supposed to be integer.

Lemma 5.2. *[Power-like potentials] Suppose that $q \leq CQ$ with $Q(t) = (1+t)^{-2\gamma}$, $\gamma > 1$.*

- (i) *Then for any $R \geq 0$*

$$(5.2) \quad M(\lambda; \pm gq, (R, \infty)) \leq C(g^{1/2}(1+R)^{1-\gamma} + 1), \quad \forall k \geq 0,$$

uniformly in $\lambda \in \bar{l}_l$ and $R \geq 0$.

- (ii) *If the condition (4.21) is satisfied, then*

$$(5.3) \quad M(\lambda; \pm gq_R) \leq Cg^{1/2}(1+R)^{1-\gamma}, \quad \forall k \geq 0,$$

uniformly in $\lambda \in I$ with a closed interval I from (4.21).

Lemma 5.3. *[Exponential potentials] Suppose that $q \leq CQ$ with $Q(t) = e^{-2\kappa t}$, $\kappa > 0$.*

- (i) *Then for any $R > 0$*

$$(5.4) \quad M(\lambda; \pm gq, (R, \infty)) \leq C(g^{1/2}e^{-\kappa R} + 1), \quad \forall k \geq 0,$$

uniformly in $\lambda \in \bar{l}_l$ and $R \geq 0$.

- (ii) *If the condition (4.21) is satisfied, then*

$$(5.5) \quad M(\lambda; \pm gq_R) \leq Cg^{1/2}e^{-\kappa R}, \quad \forall R \geq 0,$$

uniformly in $\lambda \in I$ with a closed interval I from (4.21).

Proofs of Lemmas 5.2 and 5.3. Due to the monotonicity of the function $M(\lambda; \pm V)$ in V , (see Subsect. 4.1), it is sufficient to obtain the estimates for the “model” potential Q . For a power-like Q , we have

$$\left[\int_R^{R+1} (1+t)^{-2\gamma} dt \right]^{1/2} + \int_R^\infty (1+t)^{-\gamma} dt \leq C(1+R)^{1-\gamma},$$

which implies (5.2) by virtue of (4.26). Similarly, (4.22) leads to (5.3).

The proof of Lemma 5.3 is the same. $\square$

Proof of Theorem 5.1. According to (5.3) (resp. (5.5)) the series (4.30) is convergent for $\gamma > 2$ (resp. all $\varkappa > 0$), which ensures the validity of (4.31).

In the exponential case, if $\varkappa > \ln b$, then the series (4.28) is also convergent, which leads to the asymptotics (4.29) by Theorem 4.4. $\square$

5.2. Non-Weylian asymptotics. The rest of the paper is focused on the situations when the Weyl formula fails, and the asymptotics of the counting functions (4.1)–(4.4) depends on the behaviour of the potential at infinity. We concentrate on bounded potentials q behaving like Q (see (5.1)) at infinity. The precise meaning of this phrase will be made clear later.

As in the previous section, the asymptotics of $M(\lambda; \mathbf{A}_{\pm gq})$ and $\widetilde{M}(\lambda; \mathbf{A}_{\pm gq})$ will be deduced from the asymptotics of the individual counting functions $M(\lambda; \pm gq_k)$, $k \geq 0$ for the operators for the operators $A_{\pm gq_k}$. In the case of the power-like potential q the total number of eigenvalues of $A_{\mathbf{s}q}$ in each gap may become infinite. More precisely, if $q = (1+t)^{-2\gamma}$, $\mathbf{s} = -1$ (resp. $\mathbf{s} = +1$) and $\gamma \leq 1$ then the eigenvalues accumulate at the upper (resp. lower) end of the gap $\mathfrak{l}_l$. In this connection it is convenient to introduce the notion of an *admissible point* $\lambda \in \mathfrak{l}_l$. From now on we fix the number $l \geq 0$ and denote $\mathfrak{l}_l = (\lambda_-, \lambda_+)$. If $l = 0$, then $\lambda_- = -\infty$ and $\lambda_+ = \theta^2$. If $l \geq 1$, then $\lambda_{\pm} = (\pi l \pm \theta)^2$. In the definition below, to avoid unnecessary repetitions, by $[-\infty, \lambda_+]$, $[-\infty, \lambda_+)$ we understand the intervals $(-\infty, \lambda_+]$, $(-\infty, \lambda_+)$.

Definition 5.4. Let $Q(t) = (1+t)^{-2\gamma}$. Then a point $\lambda \in [\lambda_-, \lambda_+]$ is said to be γ_0 -admissible, $\gamma_0 > 0$, if

$$\lambda \in \begin{cases} (\lambda_-, \lambda_+], & \gamma \leq \gamma_0, \mathbf{s} = +1; \\ [\lambda_-, \lambda_+), & \gamma \leq \gamma_0, \mathbf{s} = -1; \\ [\lambda_-, \lambda_+], & \gamma > \gamma_0. \end{cases}$$

For $Q(t) = e^{-2\varkappa t}$ any point $\lambda \in [\lambda_-, \lambda_+]$ is said to be γ_0 -admissible with any $\gamma_0 > 0$.

Clearly, for any two positive numbers γ_0, γ_1 , $\gamma_0 < \gamma_1$, any γ_1 -admissible λ is automatically γ_0 -admissible. For the model potential $Q(t) = (1+t)^{-2\gamma}$ the number $M(\lambda; \pm gq_n)$ is finite for all $g > 0$ if λ is 1-admissible. For the exponential model potential $Q(t) = e^{-2\varkappa t}$ the quantity $M(\lambda; \pm gq_n)$ is finite for all $\lambda \in [\lambda_-, \lambda_+]$.

5.3. Results for the functions $\widetilde{M}(\lambda; \mathbf{A}_{\mathbf{s}gq})$, $\widetilde{N}(\lambda_1, \lambda_2; \mathbf{A}_{\mathbf{s}gq})$. This subsection contains the results on the asymptotics of $\widetilde{M}(\lambda; \mathbf{A}_{\pm gq})$ and $\widetilde{N}(\lambda_1, \lambda_2; \mathbf{A}_{\pm gq})$. Their proofs require some technical preparations which we give in Sections 6, 7. The proofs are completed in Section 8. Our results for the functions (4.1), (4.2) require different techniques and are much less complete than those for $\widetilde{M}$ and $\widetilde{N}$; they are presented in Sect. 9.

Recall that in contrast to the spectrum of the “individual” operator A_0 the spectrum of $\mathbf{A}_0$ contains eigenvalues $\lambda_l = (\pi l)^2$ of infinite multiplicity. Thus, when stating the

results we assume that $\lambda, \lambda_1, \lambda_2$ satisfy (4.21) and are 2-admissible. The constants in all the estimates below are

- uniform in $\lambda, \lambda_1, \lambda_2$ varying within any closed interval I of 2-admissible points, satisfying (4.21),
- independent of the coupling constant g .

We begin with the power-like potentials.

Theorem 5.5. *Let q satisfy the condition*

$$(5.6) \quad q(t) = Q(t)(1 + o(1)), \quad t \rightarrow \infty, \quad Q(t) = (1 + t)^{-2\gamma},$$

and one of the following two conditions be fulfilled:

1. $\gamma \in (0, 2)$ and $\mathfrak{s} = -1$;
2. The exponent $\gamma > 0$ is arbitrary and $\mathfrak{s} = +1$.

Suppose that λ is 2-admissible and satisfies (4.21). Then

$$(5.7) \quad \lim_{g \rightarrow \infty} g^{-\frac{1}{\gamma}} \widetilde{M}(\lambda; \mathbf{A}_{\pm gq}) = \pm \int_0^\infty \int_0^\infty [\rho(\lambda) - \rho(\lambda \mp (s + \sigma)^{-2\gamma})] ds d\sigma,$$

where ρ is the density of states for the operator A_0 .

Remark 5.6. A simple change of variables leads to another expression for the asymptotic coefficient:

$$\lim_{g \rightarrow \infty} g^{-\frac{1}{\gamma}} \widetilde{M}(\lambda; \mathbf{A}_{\pm gq}) = \pm \int_0^\infty \int_\beta^\infty [\rho(\lambda) - \rho(\lambda \mp s^{-2\gamma})] ds d\beta.$$

In Sect. 7 we shall show that the asymptotic coefficients in the r.h.s. of (5.7) and that in Theorem 5.9 below, are finite.

Note that in contrast to $\mathfrak{s} = -1$, the above formula describes the asymptotics of $\widetilde{M}(\lambda; \mathbf{A}_{\mathfrak{s}gq})$ with $\mathfrak{s} = +1$ for all positive γ . If $\mathfrak{s} = -1$, then the case $\gamma = 2$ is critical in the sense that for $\gamma > 2$ the Weyl asymptotics is applicable instead of (5.7) (cf. Lemma 5.2). We point out however that for the *individual* counting function $M(\lambda; -gq_n)$ the critical case is $\gamma = 1$ (see Theorem 4.3).

To find a formula for $\widetilde{M}$ in the case $\gamma = 2$ we need to introduce more restrictions on q .

Condition 5.7. *Let $q \in C^1(\mathbb{R}_+)$ be a function such that*

$$\begin{aligned} cQ(t) &\leq q(t) \leq CQ(t), \quad \forall t \in \mathbb{R}_+, \\ |q'(t)| &\leq CQ(t). \end{aligned}$$

Now we are in position to study the critical case:

Theorem 5.8. *Suppose that q satisfies (5.6) with $\gamma = 2$ and Condition 5.7. Let λ be 2-admissible and satisfy (4.21). Then*

$$\lim_{g \rightarrow \infty} g^{-1/2} (\ln g)^{-1} \widetilde{M}(\lambda; \mathbf{A}_{-gq}) = (4\pi)^{-1}, \quad g \rightarrow \infty.$$

The next theorem gives an asymptotic formula for the number $\tilde{N}(\lambda_1, \lambda_2)$:

Theorem 5.9. *Suppose that q satisfies (5.6) with some $\gamma > 0$, and that in the case $\mathfrak{s} = -1$, $\alpha \geq 2$, Condition 5.7 is also fulfilled. Let λ_1, λ_2 be 2-admissible and satisfy (4.21). Then*

$$(5.8) \quad \lim_{g \rightarrow \infty} g^{-\frac{1}{\gamma}} \tilde{N}(\lambda_1, \lambda_2, \mathbf{A}_{\pm gq}) = \int_0^\infty \int_0^\infty [\rho(\lambda_2 \mp (t + \sigma)^{-2\gamma}) - \rho(\lambda_1 \mp (t + \sigma)^{-2\gamma})] dt d\sigma,$$

as $g \rightarrow \infty$.

We point out that the asymptotics of $\tilde{N}(\lambda_1, \lambda_2)$ is described by the density of states $\rho(\lambda)$ for all $\gamma > 0$. Under the conditions of Theorem 5.5 the asymptotics (5.8) can be immediately deduced from (5.7) with the help of (4.11). On the contrary, for $\alpha > 2$ and $\mathfrak{s} = -1$ the behaviour of $\tilde{N}$ can not be inferred from the asymptotics of $\tilde{M}(\lambda)$ which is given by the Weyl term, see Lemma 5.2.

Let us proceed to the exponential potentials. From Lemma 5.3 we know that for the case $q \leq CQ$, $Q(t) = e^{-2\kappa t}$, $\mathfrak{s} = -1$, the asymptotics of $\tilde{M}(\lambda; \mathbf{A}_{\mathfrak{s}q})$ is described by the Weyl formula (4.31). The next theorem gives an answer in the case $\mathfrak{s} = +1$. Below $g_0 > e$ is a constant.

Theorem 5.10. *Suppose that*

$$(5.9) \quad cQ(t) \leq q(t) \leq CQ(t), \quad \forall t \geq R_0,$$

with $Q(t) = e^{-2\kappa t}$, $\kappa > 0$ and some $R_0 \geq 0$. Let $\lambda, \lambda_1, \lambda_2$ be arbitrary numbers satisfying (4.21). Then

$$(5.10) \quad \tilde{M}(\lambda; \mathbf{A}_{gq}) = \frac{1}{8\kappa^2} \rho(\lambda) (\ln g)^2 + O(\ln g), \quad g \geq g_0,$$

and

$$(5.11) \quad \tilde{N}(\lambda_1, \lambda_2; \mathbf{A}_{gq}) \leq C \ln g, \quad g \geq g_0.$$

The next result complements the Weyl formula (4.31) by providing an estimate for the function $\tilde{N}(\lambda_1, \lambda_2; \mathbf{A}_{-gq})$:

Theorem 5.11. *Suppose that q fulfills Condition 5.7 with $Q(t) = e^{-2\kappa t}$. Let λ_1, λ_2 be arbitrary numbers satisfying (4.21). Then*

$$(5.12) \quad \tilde{N}(\lambda_1, \lambda_2; \mathbf{A}_{-gq}) \leq C (\ln g)^2, \quad g \geq g_0.$$

6. INDIVIDUAL ESTIMATES AND WEYL ASYMPTOTICS WITH A REMAINDER

6.1. Individual estimates. Here we obtain further estimates for individual counting functions $M(\lambda; \pm gq, \Delta)$. Although our ultimate objective is to establish asymptotic formulae for the counting functions $M(\lambda; \pm gq_k)$ with integer non-negative k 's, most of

the results in this section are uniform with respect to a wide class of potentials, including the shifted potentials $q_R, R \geq 0$.

Unless stated otherwise, in this section we always assume that the points $\lambda, \lambda_1, \lambda_2 \in [\lambda_-, \lambda_+]$ are 1-admissible. The constants in all the estimates obtained below are

- uniform in $\lambda, \lambda_1, \lambda_2$ varying within any closed interval $I \subset [\lambda_-, \lambda_+]$ of 1-admissible points;
- independent of the coupling constant g .

Whenever possible we treat the power-like and exponential potentials simultaneously. It is convenient to use the notation

$$(6.1) \quad \alpha = \begin{cases} g^{\frac{1}{2\gamma}}, & \text{if } Q(t) = (1+t)^{-2\gamma}; \\ \frac{1}{2\kappa} \ln g, & \text{if } Q(t) = e^{-2\kappa t}. \end{cases}$$

We always assume that $g \geq g_0 > e$, so $\alpha \geq \alpha_0$ with some $\alpha_0 > 0$ in both cases.

The following simple Lemma will be repeatedly used:

Lemma 6.1. *Suppose that $q(t) \leq CQ(t)$.*

- (i) *If $Q(t) = (1+t)^{-2\gamma}$, then for all $R \geq 1$*

$$(6.2) \quad |M(\lambda; \pm gq, \mathbb{R}_+) - M(\lambda; \pm gq, (0, R\alpha))| \leq C'\alpha.$$

with a constant C' depending only on C . Moreover,

$$(6.3) \quad \limsup_{R \rightarrow \infty} \sup_{g \geq g_0} \alpha^{-1} |M(\lambda; \pm gq, \mathbb{R}_+) - M(\lambda; \pm gq, (0, R\alpha))| = 0.$$

- (ii) *If $Q(t) = e^{-2\kappa t}$, then*

$$(6.4) \quad |M(\lambda; \pm gq, \mathbb{R}_+) - M(\lambda; \pm gq, (0, R\alpha))| \leq C',$$

with a constant C' depending only on C .

Proof. In view of the decoupling principle (4.13)

$$|M(\lambda; \pm gq, \mathbb{R}_+) - M(\lambda; \pm gq, (0, R\alpha))| \leq 2 + M(\lambda; \pm qq, (R\alpha, \infty)).$$

If $Q(t) = (1+t)^{-2\gamma}$ and $\gamma > 1$, then by (5.2) the last term in the r.h.s. does not exceed

$$g^{1/2}(\alpha R + 1)^{1-\gamma} + 1 \leq \alpha R^{1-\gamma} + 1.$$

This implies (6.2) and (6.3). If $\gamma \leq 1$, then $\lambda < \lambda_+$ (for $\mathfrak{s} = -1$) or $\lambda > \lambda_-$ (for $\mathfrak{s} = +1$). Thus one can choose $\hat{R}$ so as to ensure that $M(\lambda; \pm qq, (\hat{R}\alpha, \infty)) = 0$, since $|gq(t)| \leq g\hat{R}^{-2\gamma}\alpha^{-2\gamma} = \hat{R}^{-2\gamma}$ for all $t \geq \hat{R}\alpha$. Now (6.3) follows. To show (6.2) use the

decoupling principle and (4.26) to conclude that

$$\begin{aligned} M(\lambda; \pm gq, (R\alpha, \infty)) &\leq 2 + M(\lambda; \pm gq, (R\alpha, \hat{R}\alpha)) \\ &\leq C \left[\alpha \left[\int_R^{R+1} |t|^{-2\gamma} dt \right]^{1/2} + \alpha \int_R^{\hat{R}} |t|^{-\gamma} dt + 1 \right] + 2 \\ &\leq C(1 + \alpha). \end{aligned}$$

For the case $Q(t) = e^{-2\lambda t}$, using (5.4) we obtain the estimate:

$$M(\lambda; \pm gq, (R\alpha, \infty)) \leq C(g^{1/2}e^{-\lambda R} + 1) \leq C(g^{1/2-R/2} + 1) \leq C$$

for all $R \geq 1$, as required. $\square$

Lemma 6.2. *Let $q(t) \leq CQ(t)$.*

(i) *If $Q(t) = (1+t)^{-2\gamma}$, then*

$$(6.5) \quad M(\lambda; -gq) \leq \begin{cases} Cg^{1/2}, & \gamma > 1, \\ Cg^{1/2} \ln g, & \gamma = 1, \\ C\alpha, & \gamma < 1. \end{cases}$$

(ii) *Let either $Q(t) = (1+t)^{-2\gamma}$ with arbitrary $\gamma > 0$, or $Q(t) = e^{-2\lambda t}$. Then*

$$(6.6) \quad M(\lambda; gq) \leq C\alpha,$$

and

$$(6.7) \quad N(\lambda_1, \lambda_2; gq) \leq C\alpha.$$

Proof. (i) The estimate (6.5) for $\gamma > 1$ follows from (5.2). If $\gamma \leq 1$, then by (4.26),

$$\begin{aligned} M(\lambda; -gq, (0, \alpha)) &\leq Cg^{1/2} \left[\int_0^1 (1+t)^{-2\gamma} dt \right]^{1/2} + Cg^{1/2} \int_0^\alpha t^{-\gamma} dt + C \\ &\leq \begin{cases} Cg^{1/2} \ln \alpha, & \gamma = 1, \\ C\alpha, & \gamma < 1. \end{cases} \end{aligned}$$

By virtue of Lemma 6.1 this leads to (6.5).

(ii) It is clear from (4.9) that

$$M(\lambda; gq, (0, \alpha)) \leq N(\lambda; 0, (0, \alpha)) \leq C\alpha.$$

Now Lemma 6.1 gives (6.6). The estimate (6.7) follows from (6.6) and (4.10). $\square$

From these Lemmas we can immediately deduce the asymptotics of $M(\lambda; gq)$ with an exponential q .

Theorem 6.3. *Let q be a bounded function satisfying (5.9) with $Q(t) = e^{-2\kappa t}$. Then*

$$(6.8) \quad |M(\lambda; gq) - \rho(\lambda)\alpha| \leq C, \quad \forall \alpha \geq 1,$$

and

$$(6.9) \quad N(\lambda_1, \lambda_2; gq) \leq C, \quad \forall \alpha \geq 1.$$

Proof. The bound for $N(\lambda_1, \lambda_2; gq)$ immediately follows from (6.8) by (4.11), since $\rho(\lambda_1) = \rho(\lambda_2)$ for $\lambda_1, \lambda_2 \in [\lambda_-, \lambda_+]$.

By the decoupling principle (6.4) it suffices to study the counting functions $M(\lambda; gq, \Delta)$ with $\Delta = (0, \alpha)$. The result for the unperturbed function $N(\lambda; 0, \Delta)$ immediately follows from Theorem 3.4:

$$(6.10) \quad N(\lambda; 0, \Delta) = \rho(\lambda)\alpha + O(1), \quad \alpha \rightarrow \infty.$$

To handle the perturbed function split the interval Δ as follows:

$$\begin{aligned} \Delta &= \Delta_0 \cup \Delta_1 \cup \Delta_2, \\ \Delta_0 &= (0, R_0], \quad \Delta_1 = (R_0, R_1\alpha], \quad \Delta_2 = (R_1\alpha, \alpha), \end{aligned}$$

where $R_0 > 0$ is defined in (5.9). The number $R_1 > 0$ is found from the requirement

$$gq(t) \geq \lambda - \lambda_0, \quad \forall t \in (R_0, R_1\alpha],$$

where $\lambda_0 = \theta^2 = \inf \sigma(A_0)$. This implies that for R_1 we can take

$$(6.11) \quad R_1 = 1 - \frac{C'}{2\kappa\alpha},$$

with a sufficiently large $C' = C'(\lambda) > 0$. It follows from (6.11) and Theorem 3.4 that

$$N(\lambda; gq, \Delta_2) \leq N(\lambda; 0, \Delta_2) \leq (1 - R_1)\alpha + C' \leq C'', \quad \forall \alpha \geq 1,$$

Since $N(\lambda; gq, \Delta_0) \leq N(\lambda; 0, \Delta_0) \leq C$ and $N(\lambda; gq, \Delta_1) = 0$, by the decoupling principle (4.12) we have

$$N(\lambda; gq, \Delta) \leq C, \quad \forall \alpha \geq 1.$$

Now it follows from (6.10) and (4.9) that

$$|M(\lambda; gq, \Delta) - \rho(\lambda)\alpha| \leq C, \quad \alpha \geq 1,$$

which implies (6.8). □

6.2. Individual asymptotics of the Weyl type with a remainder. Even if a potential $V \leq 0$ satisfies the conditions of Theorem 4.3, the formula (4.27) fails to provide an asymptotics for $N(\lambda_1, \lambda_2; V)$ as the leading term in (4.27) does not depend on λ . Below we establish, under certain conditions on q , a Weyl-type asymptotics for $M(\lambda; -gq, \Delta)$ with a remainder, which allows us to obtain bounds on the growth of $N(\lambda_1, \lambda_2; -gq)$ as $g \rightarrow \infty$.

We begin with an asymptotic formula for the Schrödinger operator $-d^2/dt^2 - q$ on a bounded interval $\Delta \subset \mathbb{R}_+$ with the Dirichlet boundary conditions but *without* any

matching conditions. Denote the counting function of this operator by $\#(\lambda; -q, \Delta)$. The next theorem is a minor modification of a similar statement from [19]:

Theorem 6.4. *Let $q \in C^1([0, \infty))$ be a non-negative function, and let $\Delta = (R_1, R_2)$ with $0 \leq R_1 \leq R_2 < \infty$. Then for any $\lambda \in \mathbb{R}$ one has*

$$(6.12) \quad \left| \#(\lambda; -q; \Delta) - \frac{1}{\pi} \int_{\Delta} \sqrt{q(t)} \, dt \right| \leq \int_{\Delta} \frac{|q'(t)|}{4\pi(q(t) + |\lambda|)} \, dt + \frac{3\sqrt{|\lambda| + 1}}{\pi} |\Delta| + 1,$$

where the constant C does not depend on Δ, g and is uniform in λ on a compact interval.

Proof. Assume without loss of generality that $R_1 = 0$. The idea is to use the fact that the number $\#(\lambda) = \#(\lambda; -q, \Delta)$ equals the number of roots of the solution u of the equation

$$(6.13) \quad -u'' - qu = \lambda u, \quad u(0) = 0, \quad u'(0) = 1,$$

lying strictly inside Δ . To find the number of roots, represent u in the polar form:

$$(6.14) \quad u(t) = \beta(t) \sin \xi(t), \quad u'(t) = \beta(t) f(t) \cos \xi(t), \quad f = \sqrt{q + \lambda_0},$$

with a $\lambda_0 > 0$. The equation (6.13) and the above equalities define the real-valued amplitude β and the phase ξ uniquely under the assumption that ξ is continuous. Substituting (6.14) in the equation (6.13), one obtains the following non-linear equation for ξ :

$$(6.15) \quad \xi' = f + \frac{f'}{2f} \sin(2\xi) + \frac{\lambda - \lambda_0}{f} \sin^2 \xi, \quad \xi(0) = 0,$$

and a linear equation for β :

$$\beta' = -\frac{1}{f} ((\lambda - \lambda_0) \sin \xi + f' \cos \xi) \cos \xi \, \beta, \quad \beta(0) = \frac{1}{f(0)}.$$

Since β never vanishes, the number of roots of u equals the number of points $t \in \Delta$ where $\xi(t) = 0 \pmod{\pi}$. From (6.15) it is clear that $\xi'(t) = f(t) > 0$ for those t , so that

$$\pi^{-1} \xi(R_2) - 1 \leq \#(\lambda) \leq \pi^{-1} \xi(R_2).$$

Therefore (6.15) implies that

$$\left| \#(\lambda) - \frac{1}{\pi} \int_{\Delta} f \, dt \right| \leq \int_{\Delta} \frac{|f'|}{2\pi f} \, dt + \int_{\Delta} \frac{|\lambda - \lambda_0|}{\pi f} \, dt + 1.$$

Since $f \geq \sqrt{\lambda_0}$, $f' = q'(2f)^{-1}$, and $\sqrt{q + \lambda_0} - \sqrt{q} \leq \sqrt{\lambda_0}$, this leads to

$$\left| \#(\lambda) - \frac{1}{\pi} \int_{\Delta} \sqrt{q} \, dt \right| \leq \int_{\Delta} \frac{|q'|}{4\pi(q + \lambda_0)} \, dt + \int_{\Delta} \frac{|\lambda| + 2\lambda_0}{\pi \sqrt{\lambda_0}} \, dt + 1.$$

It remains to take $\lambda_0 = |\lambda| + 1$. □

Theorem 6.5. *Suppose that q satisfies Condition 5.7 and let $\Delta = (R_1, R_2)$ with $0 \leq R_1 \leq R_2 < \infty$. Then for any $\lambda \in \mathbb{R}$ one has*

$$(6.16) \quad \left| N(\lambda; -gq; \Delta) - \frac{\sqrt{g}}{\pi} \int_{\Delta} \sqrt{q(t)} dt \right| \leq C(|\Delta| + 1),$$

for all $g \geq g_0$, where the constant C does not depend on Δ, g and is uniform in λ on a compact interval.

Proof. Let us split Δ as follows:

$$\begin{aligned} \Delta &= \cup_{k=l}^m \Delta_k, \quad l = [R_1], m = [R_2], \\ \Delta_l &= (R_1, l + 1], \quad \Delta_m = (m, R_2), \\ \Delta_k &= (k, k + 1], \quad k = l + 1, \dots, m - 1. \end{aligned}$$

Then by the decoupling principle

$$|N(\lambda; \Delta) - \sum_k N(\lambda; \Delta_k)| \leq 2(m - l) \leq 2(|\Delta| + 2).$$

To study each Δ_k we use Theorem 6.4. Namely, since $|q'| \leq Q$ and $q \geq cQ$, the estimate (6.12) yields:

$$\left| N(\lambda; \Delta_k) - \frac{\sqrt{g}}{\pi} \int_{\Delta_k} \sqrt{q(t)} dt \right| \leq C$$

with a constant independent of k . Adding up these inequalities over $k = l, l + 1, \dots, m$, we arrive at (6.16). $\square$

Let us derive from this theorem the asymptotics for the counting function $M(\lambda; -gq, \Delta)$ with $\Delta = (k, \infty)$, $k > 0$.

Theorem 6.6. *Suppose that Condition 5.7 is satisfied and let $\Delta = (k, \infty)$, $k \in \mathbb{N} \cup \{0\}$. Then*

$$(6.17) \quad \left| M(\lambda; -gq, \Delta) - \frac{\sqrt{g}}{\pi} \int_k^{k+\alpha} \sqrt{q(t)} dt \right| \leq C\alpha, \quad \forall g \geq g_0,$$

uniformly in k , and

$$(6.18) \quad N(\lambda_1, \lambda_2; -gq, \Delta) \leq C\alpha, \quad \forall g \geq g_0.$$

If $Q(t) = (1 + t)^{-2\gamma}$, $\gamma > 1$ or $Q(t) = e^{-2\lambda t}$, then

$$(6.19) \quad \left| M(\lambda; -gq, \Delta) - \frac{\sqrt{g}}{\pi} \int_k^\infty \sqrt{q(t)} dt \right| \leq C\alpha, \quad \forall g \geq g_0.$$

Proof. The estimate (6.18) follows from (6.17) by virtue of (4.10).

Let us prove (6.17). Let $\Delta_1 = (k, k + \alpha]$ and $\Delta_2 = (k + \alpha, \infty)$. In view of Lemma 6.1 and the relation (4.5) it suffices to show that the distribution function $M(\lambda; -gq, \Delta_1)$ satisfies (6.17). This is a direct consequence of Theorem 6.5 and the identity (4.9).

The formula (6.19) follows from (6.17) in view of the inequality

$$\sqrt{g} \int_{k+\alpha}^{\infty} \sqrt{q(t)} dt \leq C\alpha.$$

□

7. INDIVIDUAL ASYMPTOTICS. POWER-LIKE POTENTIALS

In this section we study the individual counting function $M(\lambda; \mathbf{s} g q_k)$ for q satisfying (5.6) with $Q(t) = (1+t)^{-2\gamma}$, where $\gamma > 0$ is arbitrary for both cases $\mathbf{s} = \pm 1$. In contrast to the previous section here we focus on the asymptotics of this function under the assumption that g and k tend to infinity in a coordinated way (see Lemma 7.3 below). We shall use the following notation:

$$(7.1) \quad \alpha = g^{\frac{1}{2\gamma}}, \quad \beta = \beta_k(g) = \frac{k+1}{\alpha}.$$

Emphasise again that the main difference with the asymptotics obtained in Theorem 4.3 is that now it is determined by the density of states for the unperturbed operator A_0 .

We begin with the study of asymptotic coefficients.

7.1. Asymptotic coefficients. Introduce the asymptotic coefficients for $M(\lambda; \pm g q_k)$:

$$(7.2) \quad F_{\pm}(\sigma, \lambda) = \pm \int_0^{\infty} [\rho(\lambda) - \rho(\lambda \mp (s + \sigma)^{-2\gamma})] ds$$

and for $N(\lambda_1, \lambda_2; \pm g q_k)$:

$$(7.3) \quad \begin{aligned} G_{\pm}(\sigma, \lambda_1, \lambda_2) &= \pm (F_{\pm}(\sigma, \lambda_1) - F_{\pm}(\sigma, \lambda_2)) \\ &= \int_0^{\infty} [\rho(\lambda_2 \mp (s + \sigma)^{-2\gamma}) - \rho(\lambda_1 \mp (s + \sigma)^{-2\gamma})] ds. \end{aligned}$$

It is clear that $F_{\pm} \geq 0$ and $G_{\pm} \geq 0$ if $\lambda_1 \leq \lambda_2$. Some other useful properties of $F_{\pm}, G_{\pm}$ are collected in the next Lemma:

Lemma 7.1. *Let $\lambda, \lambda_1, \lambda_2 \in [\lambda_-, \lambda_+]$ be 1-admissible numbers. Then the integral $F_{\pm}(\sigma, \lambda)$ is finite for all $\sigma > 0$. Moreover,*

(i) *If $\gamma > 1$, then*

$$F_{-}(\sigma, \lambda) \leq C\sigma^{1-\gamma},$$

$$F_{+}(\sigma, \lambda) \leq C(1 + \sigma)^{1-\gamma},$$

for all $\sigma > 0$.

(ii) *If $\gamma \leq 1$, then $F_{+}(\sigma, \lambda) \leq C$ and*

$$F_{-}(\sigma, \lambda) \leq \begin{cases} C, & \gamma < 1, \\ C \ln(\sigma^{-1} + 1), & \gamma = 1. \end{cases}$$

(iii) If $\mathfrak{s} = +1$ and $\lambda > \lambda_-$ or $\mathfrak{s} = -1$ and $\lambda < \lambda_+$, then $F_{\pm}(\sigma, \lambda) = 0$ for all $\sigma \geq \sigma_{\pm}(\lambda) = d_{\pm}^{-\frac{1}{2\gamma}}$ with $d_{\pm} = |\lambda - \lambda_{\mp}|$, and

$$F_{\pm}(\sigma, \lambda) \geq C(\sigma_{\pm} - \sigma)^{\frac{3}{2}}, \quad \sigma \leq \sigma_{\pm}.$$

(iv) The integral

$$(7.4) \quad F_{\pm}^R(\sigma, \lambda) = \pm \int_0^R [\rho(\lambda) - \rho(\lambda \mp (s + \sigma)^{-2\gamma})] ds, \quad R > 0,$$

tends to $F_{\pm}(\sigma, \lambda)$ as $R \rightarrow \infty$ uniformly in $\sigma > 0$.

(v) For all $\gamma > 0$ one has $G_{\pm}(\sigma, \lambda_1, \lambda_2) \leq C$, $\forall \sigma > 0$.

Proof. (i) By (3.24), the integrand in the definition (7.2) does not exceed $C(s + \sigma)^{-\gamma}$ for $\mathfrak{s} = -1$ and $\min\{\rho(\lambda), C(s + \sigma)^{-\gamma}\}$ for $\mathfrak{s} = +1$. The required estimates follow immediately.

(ii) Let $\gamma \leq 1$. Let first $\mathfrak{s} = +1$, so that $\lambda \in (\lambda_-, \lambda_+]$. Define $R > 0$ to be the number such that $\lambda - R^{-2\gamma} = \lambda_-$. Consequently, $\lambda - (s + \sigma)^{-2\gamma} \geq \lambda_-$ for $s + \sigma \geq R$, and hence

$$\rho(\lambda) - \rho(\lambda - (s + \sigma)^{-2\gamma}) = 0, \quad \forall s : s + \sigma \geq R.$$

This implies that

$$F_+(\sigma, \lambda) \leq \int_0^R \rho(\lambda) ds = R\rho(\lambda) \leq C'(\lambda).$$

Consider now the case $\mathfrak{s} = -1$, so that $\lambda \in [\lambda_-, \lambda_+)$. Let $R > 0$ be the number such that $\lambda + R^{-2\gamma} = \lambda_+$. Consequently, $\lambda + (s + \sigma)^{-2\gamma} \leq \lambda_+$ for $s + \sigma \geq R$, and hence

$$\rho(\lambda + (s + \sigma)^{-2\gamma}) - \rho(\lambda) = 0, \quad \forall s : s + \sigma \geq R.$$

If $\sigma \geq R$, then $F_-(\sigma, \lambda) = 0$. If $\sigma < R$, then

$$F_-(\sigma, \lambda) \leq C \int_0^{R-\sigma} (\sigma + s)^{-\gamma} ds \leq \begin{cases} C(R), & \gamma < 1, \\ C(R) \ln(\sigma^{-1} + 1), & \gamma = 1. \end{cases}$$

(iii) For brevity consider only the case $\mathfrak{s} = -1$, so that $\lambda < \lambda_+$. For $s \geq \sigma_- - \sigma$, $\sigma_- = \sigma_-(\lambda)$, the integrand in (7.2) equals zero. Besides, as $\rho(\lambda) = \rho(\lambda_+)$, in view of (3.23) we have

$$\rho(\lambda + (s + \sigma)^{-2\gamma}) - \rho(\lambda) \geq c((s + \sigma)^{-2\gamma} - d_-)^{\frac{1}{2}} \geq c'(\sigma_- - \sigma - s)^{\frac{1}{2}},$$

$$\forall s \in (0, \sigma_- - \sigma).$$

Integrating this inequality in s , we obtain the required lower bound for $F_-(\sigma, \lambda)$. The analogous bound for $F_+(\sigma, \lambda)$ is obtained in the same way.

(iv) It suffices to notice that for any 1-admissible λ and any $R > 0$ one has

$$\pm \int_R^{\infty} [\rho(\lambda) - \rho(\lambda \mp (\sigma + s)^{-2\gamma})] ds \leq \pm \int_R^{\infty} [\rho(\lambda) - \rho(\lambda \mp s^{-2\gamma})] ds$$

for all $\sigma \geq 0$.

(v) Arguing as on the previous step, it suffices to prove that the integral of the form (7.3) over a finite interval is bounded uniformly in $\sigma > 0$. By (3.24)

$$|\rho(\lambda_2 \mp (s + \sigma)^{-2\gamma}) - \rho(\lambda_1 \mp (s + \sigma)^{-2\gamma})| \leq C|\lambda_1 - \lambda_2|^{1/2},$$

which provides the required boundedness. $\square$

When studying the sum of the counting functions, we shall need some properties of the sum of asymptotic coefficients $F_{\pm}(\beta_k, \lambda)$:

Lemma 7.2. (i) *Suppose that λ is a 1-admissible number and $\delta = \delta(\alpha)$ is a bounded function such that*

$$(7.5) \quad \delta \leq 1, \quad \alpha \delta^{2\gamma+1} \rightarrow \infty, \quad \text{as } \alpha \rightarrow \infty.$$

Then for any fixed $A \geq \sup_{\alpha} \delta(\alpha)$ one has

$$(7.6) \quad \lim \left[\alpha^{-1} \sum_{[\delta\alpha] \leq k \leq [A\alpha]} F_{\pm}(\beta_k, \lambda) - \int_{\delta}^A F_{\pm}(\sigma, \lambda) d\sigma \right] = 0, \quad g \rightarrow \infty.$$

(ii) *If $\lambda, \lambda_1, \lambda_2$ are 2-admissible, then the integrals*

$$\int_0^{\infty} F_+(\sigma, \lambda) d\sigma, \quad \int_0^{\infty} G_{\pm}(\sigma, \lambda_1, \lambda_2) d\sigma$$

are finite. If in addition $\gamma \in (0, 2)$, then the integral

$$\int_0^{\infty} F_-(\sigma, \lambda) d\sigma$$

is also finite.

Proof. For brevity we omit λ from the notation of $F_{\pm}$.

(i) Let $\sigma \in (k, k+1]$ be an arbitrary number, and let

$$\lambda_1 = \lambda \pm (s + \beta_k)^{-2\gamma}, \quad \lambda_2 = \lambda \pm (s + \sigma/\alpha)^{-2\gamma}, \quad t > 0.$$

Observe that

$$|\lambda_1 - \lambda_2| \leq 2\gamma\delta^{-2\gamma-1}\alpha^{-1}.$$

Now it follows from (3.24) that

$$|\rho(\lambda_1) - \rho(\lambda_2)| \leq CE_1(\alpha, \delta), \quad E_1(\alpha, \delta) = \alpha^{-1/2}\delta^{-\gamma-1/2}.$$

Thus, by definition (7.4), for each $R > 0$ one has

$$\left| F_{\pm}^R(\beta_k) - \int_k^{k+1} F_{\pm}^R(\sigma/\alpha) d\sigma \right| \leq CRE_1(\alpha, \delta),$$

or, changing the variable under the integral,

$$\left| F_{\pm}^R(\beta_k) - \alpha \int_{k/\alpha}^{(k+1)/\alpha} F_{\pm}^R(\sigma) d\sigma \right| \leq CRE_1(\alpha, \delta).$$

Therefore

$$\begin{aligned} \left| \alpha^{-1} \sum_{k=[\delta\alpha]}^{[A\alpha]} F_{\pm}^R(\beta_k) - \int_{\delta}^A F_{\pm}^R(\sigma) d\sigma \right| \\ \leq CARE_1(\alpha, \delta) + \int_{[\delta\alpha]\alpha^{-1}}^{\delta} F_{\pm}^R(\sigma) d\sigma + \int_A^{([A\alpha]+1)\alpha^{-1}} F_{\pm}^R(\sigma) d\sigma. \end{aligned}$$

Clearly, the first term tends to zero under the conditions (7.5). The last two integrals tend to zero as $\alpha \rightarrow \infty$ by Lemma 7.1(i), (ii). Consequently, for each $R > 0$

$$\begin{aligned} \limsup \left| \alpha^{-1} \sum_{k=[\delta\alpha]}^{[A\alpha]} F_{\pm}(\beta_k) - \int_{\delta}^A F_{\pm}(\sigma) d\sigma \right| \\ \leq A \limsup \max_{[\delta\alpha] \leq k \leq [A\alpha]} |F_{\pm}^R(\beta_k) - F_{\pm}(\beta_k)| \\ + \int_{\delta}^A |F_{\pm}^R(\sigma) - F_{\pm}(\sigma)| d\sigma, \end{aligned}$$

where $\limsup$ is taken under the conditions (7.5). Recall that by Lemma 7.1 $F_{\pm}^R(\sigma)$ converges to $F_{\pm}(\sigma)$ as $R \rightarrow \infty$ uniformly in $\sigma > 0$. Thus the r.h.s. of the above inequality vanishes as $R \rightarrow \infty$. This proves (7.6).

(ii) By Lemma 7.1(i), (v), and also by definition (7.3), the functions F_{+} and $G_{\pm}$ are integrable in σ for $\gamma > 2$.

If $\gamma < 2$, then $F_{\pm}, G_{\pm}$ have compact support due to Lemma 7.1(iii). They are also integrable near $\sigma = 0$ by virtue of Lemma 7.1(i).

If $\gamma = 2$, then the same applies to F_{+} and $G_{\pm}$ again by Lemma 7.1(i), (iii), (v). $\square$

Lemma 7.2 guarantees that the asymptotic coefficients in Theorems 5.5 and 5.9 are finite.

7.2. Asymptotics of $M(\lambda; \pm gq_k)$.

Lemma 7.3. *Let q satisfy (5.6), and let $\lambda, \lambda_1, \lambda_2 \in [\lambda_-, \lambda_+]$ be 1-admissible. Then for any function $\delta = \delta(\alpha)$ satisfying (7.5) and any fixed $A > \sup_{\alpha} \delta(\alpha)$, one has*

$$(7.7) \quad \lim_{\delta \leq \beta_k \leq A} \max |\alpha^{-1} M(\lambda; \pm gq_k) - F_{\pm}(\beta_k(g), \lambda)| = 0,$$

$$(7.8) \quad \lim_{\delta \leq \beta_k \leq A} \max |\alpha^{-1} N(\lambda_1, \lambda_2; \pm gq_k) - G_{\pm}(\beta_k(g), \lambda_1, \lambda_2)| = 0,$$

as $g \rightarrow \infty$.

Proof. Note without further ado, that (7.8) is a direct consequence of (7.7) in view of (4.11).

Let us concentrate on the proof of (7.7). By (6.3) it suffices to establish the required asymptotic formula for the counting function $M(\lambda; \pm gq_k, \Delta)$ with $\Delta(g) = (0, R\alpha]$ and

afterwards take R to infinity. The proof of this fact is an adaptation of the corresponding argument from [19].

Suppose first that $q(t) = Q(t)$. Let us split $(0, R]$ into L identical subintervals $(s_{j-1}, s_j]$, $j = 1, 2, \dots, L$, so that $s_0 = 0$, $s_L = R$ and $s_{j+1} - s_j = RL^{-1}$, and denote

$$\Delta_j = (s_{j-1}\alpha, s_j\alpha], \quad j = 1, 2, \dots, L.$$

Define step functions $q_{n,1}, q_{n,2}$:

$$\begin{aligned} q_{k,1}(t) &= q_k(s_{j-1}\alpha) = \frac{1}{g(s_{j-1} + \beta)^{2\gamma}}, \quad t \in \Delta_j, \\ q_{k,2}(t) &= q_k(s_j\alpha) = \frac{1}{g(s_j + \beta)^{2\gamma}}, \quad t \in \Delta_j. \end{aligned}$$

Here we have denoted $\beta = \beta_k$. Further proof is for the case $\mathbf{s} = -1$ only. The other case is done in the same way. Clearly, $q_{k,2} \leq q_k \leq q_{k,1}$, and hence the counting function of the operator $A_0 - gq_k$ with the Dirichlet conditions at the ends of the interval Δ satisfies the two-sided estimate

$$N(\lambda; -gq_{k,2}, \Delta) \leq N(\lambda; -gq_k, \Delta) \leq N(\lambda; -gq_{k,1}, \Delta).$$

Let us find the asymptotics of the r.h.s. We are going to use the decoupling principle again:

$$(7.9) \quad N(\lambda; -gq_{k,1}, \Delta) \leq \sum_{j=1}^L N(\lambda; -gq_{k,1}, \Delta_j) + 2(L-1).$$

Now, using Theorem 3.4, we get for each j

$$\begin{aligned} \alpha^{-1}N(\lambda; -gq_{k,1}, \Delta_j) &= (s_j - s_{j-1})|\Delta_j|^{-1}N(\lambda + (s_{j-1} + \beta)^{-2\gamma}; 0, \Delta_j) \\ &\leq (s_j - s_{j-1})\rho(\lambda + (s_{j-1} + \beta)^{-2\gamma}) \\ &\quad + C(1 + \sqrt{|\lambda| + (s_{j-1} + \beta)^{-2\gamma}})\alpha^{-1}, \end{aligned}$$

with a universal constant C . Since $\beta \geq \delta$, we obtain from (7.9) that

$$\begin{aligned} \alpha^{-1}N(\lambda; -gq_{k,1}, \Delta) - \sum_j (s_j - s_{j-1})\rho(\lambda + (s_{j-1} + \beta)^{-2\gamma}) \\ \leq (L + C(\lambda) + C\delta^{-\gamma})\alpha^{-1}. \end{aligned}$$

This can be rewritten as

$$\begin{aligned} (7.10) \quad \alpha^{-1}N(\lambda; -gq_{k,1}, \Delta) - \sum_j \int_{\Delta_j} \rho(\lambda + (s_{j-1} + \beta)^{-2\gamma}) ds \\ \leq (L + C(\lambda) + C\delta^{-\gamma})\alpha^{-1}. \end{aligned}$$

To replace the sum in the l.h.s. by the integral, we use the Hölder property (3.24) with

$$\lambda_1 = \lambda + (s_{j-1} + \beta)^{-2\gamma}, \quad \lambda_2 = \lambda + (t + \beta)^{-2\gamma}.$$

Observe that

$$\begin{aligned} |\lambda_1 - \lambda_2| &= |(s_{j-1} + \beta)^{-2\gamma} - (t + \beta)^{-2\gamma}| \\ &\leq 2\gamma\delta^{-2\gamma-1}|s_{j-1} - t| \leq 2\gamma R\delta^{-2\gamma-1}L^{-1}, \quad \forall t \in \Delta_j. \end{aligned}$$

Now we infer from (3.24) that

$$\left| \rho(\lambda + (s_{j-1} + \beta)^{-2\gamma}) - \rho(\lambda + (t + \beta)^{-2\gamma}) \right| \leq C\delta^{-\gamma-1/2}R^{1/2}L^{-1/2}, \quad t \in \Delta_j.$$

Substituting this estimate into (7.10), we get

$$\begin{aligned} \alpha^{-1}N(\lambda; -gq_{k,1}, \Delta) - \int_0^R \rho(\lambda + (s + \beta)^{-2\gamma})ds &\leq CE(\alpha, \delta, L; R), \\ E(\alpha, \delta, L; R) &= (L + 1 + \delta^{-\gamma})\alpha^{-1} + \delta^{-\gamma-1/2}R^{3/2}L^{-1/2}. \end{aligned}$$

Arguing similarly, we arrive at the analogous lower bound for $N(\lambda; -gq_{k,2}, \Delta)$. Consequently,

$$\left| \alpha^{-1}N(\lambda; -gq_k, \Delta) - \int_0^R \rho(\lambda + (s + \beta)^{-2\gamma})ds \right| \leq CE(\alpha, \delta, L; R)$$

In view of Theorem 3.4 we also have

$$\left| \alpha^{-1}N(\lambda; 0, \Delta) - \int_0^R \rho(\lambda)ds \right| \leq C\alpha^{-1}.$$

By (4.9), in combination with the previous estimate this gives

$$\left| \alpha^{-1}M(\lambda; -gq_k, \Delta) - F_-^R(\beta_k, \lambda) \right| \leq CE(\alpha, \delta, L; R)$$

The parameter L can be chosen so as to insure that $E \rightarrow 0$ as $\alpha \rightarrow \infty$. Indeed, in view of (7.5) $\alpha\delta^{2\gamma+1} \rightarrow \infty$ as $\alpha \rightarrow \infty$. Therefore, defining

$$L = [\alpha^{1/2}\delta^{-\gamma-1/2}],$$

we guarantee that

$$\delta^{-2\gamma-1}L^{-1} \sim \alpha^{-1/2}\delta^{-\gamma-1/2} \rightarrow 0, \quad L\alpha^{-1} \sim \alpha^{-1/2}\delta^{-\gamma-1/2} \rightarrow 0, \quad \alpha \rightarrow \infty.$$

Taking R to infinity and referring to Lemma 7.1(iv), we obtain (7.7), thus completing the proof for $q(t) = Q(t)$.

It remains to include the potentials satisfying (5.6). To this end note that under the condition (5.6), for any $\varepsilon > 0$

$$Q_k(t)(1 - \varepsilon) \leq q_k(t) \leq Q_k(t)(1 + \varepsilon),$$

if k is sufficiently large. Thus, using the monotonicity of $M(\lambda; V)$ in V (see Sect. 4) and the asymptotics (7.7) for $q = Q$ we easily deduce (7.7) for the general case. $\square$

In conclusion note that we shall not need the asymptotics (7.8) in what follows.

8. ASYMPTOTICS OF $\widetilde{M}(\lambda, \mathbf{A}_{\pm gq})$: PROOF OF THEOREMS 5.5, 5.8, 5.9, 5.10, 5.11

Throughout this section we assume that $\lambda, \lambda_1, \lambda_2$ are 2-admissible and satisfy (4.21).

8.1. Proof of Theorems 5.5 and 5.9. Recall that in Theorem 5.5 we assume that either $\mathbf{s} = -1$, $\gamma < 2$, or $\mathbf{s} = +1$ and $\gamma > 0$ is arbitrary. In Theorem 5.9 $\gamma > 0$ is arbitrary, but if $\mathbf{s} = -1$ and $\gamma \geq 2$, then the potential q satisfies Condition 5.7.

Step I. To begin with we show that “small” or “large” values of k do not contribute to the sums $\widetilde{M}$ and $\widetilde{N}$.

Suppose that $k \geq A\alpha$ with some fixed $A > 0$. Then for $\gamma \leq 2$ and large A the perturbation $gq_k \leq CA^{-2\gamma}$ is small and therefore $M(\lambda, \pm gq_k) = 0$, since λ is strictly inside the gap. For $\gamma > 2$, by (5.3) we have

$$\sum_{k \geq A\alpha} M(\lambda; \pm gq_k) \leq Cg^{1/2}(A\alpha)^{2-\gamma} = C'A^{2-\gamma}\alpha^2.$$

By (4.11) a similar bound holds for the sum of the functions $N(\lambda_1, \lambda_2; \pm gq_k)$ for all $\gamma > 0$. These calculations again show that $k \geq A\alpha$ do not contribute as A grows.

Suppose that $k \leq \delta\alpha$. If $\gamma < 2$ and $\mathbf{s} = -1$, then the bounds (5.3) (for $\gamma > 1$) and (6.5) (for $\gamma \leq 1$) ensure that for $\delta > 0$

$$(8.1) \quad \sum_{k \leq \delta\alpha} M(\lambda; -gq_k) \leq \begin{cases} C\delta^{2-\gamma}\alpha^2, & 1 < \gamma < 2; \\ C\delta\alpha^2 \ln \alpha, & \gamma = 1; \\ C\delta\alpha^2, & \gamma < 1. \end{cases}$$

This means that the share of this sum becomes small when $\delta \rightarrow 0$. For $\gamma \neq 1$ we can take δ to be arbitrarily small constant, *independent of* α . With $\gamma = 1$ we must be more careful. Since we want to obtain the asymptotics of order α^2 , we should “kill” the “ln” term in the estimate by choosing δ to be dependent on α , but in a very mild way: $\delta = \alpha^{-\eta}$ with a parameter $\eta < (1 + 2\gamma)^{-1}$, so that the condition (7.5) from Lemma 7.3 is satisfied. For $\mathbf{s} = +1$ the estimate (6.6) yields:

$$\sum_{k \leq \delta\alpha} M(\lambda; gq_k) \leq C_\gamma \delta \alpha^2, \quad \forall \gamma > 0.$$

As in the case $\mathbf{s} = -1$ and $\gamma \neq 1$, it is possible to take δ to be an arbitrarily small constant. However, for the sake of uniformity, we take $\delta = \alpha^{-\eta}$, for both signs $\mathbf{s} = \pm 1$. Consequently,

$$\sum_{k \leq \delta\alpha} M(\lambda; \pm gq_k) = o(\alpha^2)$$

and the condition (7.5) is satisfied. In the case of the function $\tilde{N}$ the estimate (6.7) guarantees that

$$(8.2) \quad \sum_{k \leq \delta\alpha} N(\lambda_1, \lambda_2; \mathfrak{s} g q_k) \leq C_\gamma \delta \alpha^2 = o(\alpha^2),$$

for $\mathfrak{s} = +1$ and all $\gamma > 0$. If $\mathfrak{s} = -1$, $\gamma < 2$, then the same bound follows from (8.1) and (4.11). In the case $\mathfrak{s} = -1$, $\gamma \geq 2$ the estimate (8.2) is a direct consequence of (6.18). Thus, it remains to study the sums (4.3), (4.4) only over the numbers

$$[\delta\alpha] \leq k \leq [A\alpha],$$

with $\delta = \alpha^{-\eta}$, $\eta < (1 + 2\gamma)^{-1}$, and a fixed $A \geq \sup_\alpha \delta$.

Step 2. We use the notation (7.1). Estimate using (7.6):

$$\begin{aligned} \limsup \left| \alpha^{-2} \sum_{k=[\delta\alpha]}^{[A\alpha]} M(\lambda; \pm g q_k) - \int_{\delta}^A F_{\pm}(\sigma, \lambda) d\sigma \right| \\ \leq \limsup \alpha^{-1} \sum_{k=[\delta\alpha]}^{[A\alpha]} \left| \alpha^{-1} M(\lambda; \pm g q_k) - F_{\pm}(\beta_k, \lambda) \right| \\ \leq A \limsup \max_{[\delta\alpha] \leq k \leq [A\alpha]} \left| \alpha^{-1} M(\lambda; \pm g q_k) - F_{\pm}(\beta_k, \lambda) \right|. \end{aligned}$$

The r.h.s. tends to zero by Lemma 7.3. Consequently

$$\lim \left[\alpha^{-2} \sum_{k=[\delta\alpha]}^{[A\alpha]} M(\lambda; \pm g q_k) - \int_{\delta}^A F_{\pm}(\sigma, \lambda) d\sigma \right] = 0, \quad g \rightarrow \infty.$$

By (4.11) and (7.3) this equality implies that

$$\lim \left[\alpha^{-2} \sum_{k=[\delta\alpha]}^{[A\alpha]} N(\lambda_1, \lambda_2; \pm g q_k) - \int_{\delta}^A G_{\pm}(\sigma, \lambda_1, \lambda_2) d\sigma \right] = 0, \quad g \rightarrow \infty.$$

Referring to Step I of the proof and Lemma 7.2(ii), we can now replace the lower and upper limits of summation and integration by 0 and ∞ respectively. This completes the proof of Theorems 5.5, 5.9. $\square$

8.2. Proof of Theorem 5.8. By (5.6),

$$\int_0^\infty \sqrt{q_k(t)} dt = \int_0^\infty \frac{1 + o(1)}{(1 + k + t)^2} dt = \frac{1 + \epsilon_k}{k + 1},$$

where $\epsilon_k \rightarrow 0$ as $k \rightarrow \infty$. From here we find by virtue of Theorem 6.6 that the function $M((\lambda))$ has the following asymptotics:

$$(8.3) \quad \left| M(\lambda; -g q_k) - \frac{g^{1/2}}{\pi(k + 1)} \right| \leq C g^{1/4} + C' g^{1/2} \frac{\epsilon_k}{k + 1},$$

uniformly in $k \geq 0$. Observe that the components with numbers $k \geq Ag^{1/4}$ do not contribute if A is sufficiently large, since $|gq_k(t)| \leq CA^{-4}$. Let us turn to the remaining terms:

$$\left| \widetilde{M}(\lambda, \mathbf{A}_{-gq}) - \frac{g^{1/2} \ln g}{4\pi} \right| \leq \sum_{k \leq Ag^{1/4}} |M(\lambda; -gq_k) - g^{1/2}(k+1)^{-1}\pi^{-1}| \\ + g^{1/2}\pi^{-1} \left| \sum_{k \leq Ag^{1/4}} (1+k)^{-1} - \ln g/4 \right|.$$

The second term in the r.h.s. is of order $O(g^{1/2})$ in view of the known formula for the partial sum of the harmonic series. By (8.3) the first term is bounded by

$$CAg^{1/2} + C'g^{1/2} \sum_{k \leq Ag^{1/4}} \frac{\epsilon_k}{1+k}.$$

Since $\epsilon_k \rightarrow 0$ as $k \rightarrow \infty$, this quantity is of order $o(g^{1/2} \ln g)$. $\square$

8.3. Proof of Theorems 5.10, 5.11. Rewrite the sum $\widetilde{M}$ in the form

$$\widetilde{M}(\lambda, \mathbf{A}_{gq}) = \sum_k M(\lambda; ge^{-2\kappa k} \hat{q}^{(k)}), \quad \hat{q}^{(k)} = e^{2\kappa k} q_k.$$

Since $\hat{q}^{(k)}, k \geq 0$, satisfies the bound (5.9) for all $t \geq R_0$, from Theorem 6.3 we obtain that

$$|M(\lambda; ge^{-2\kappa k} \hat{q}^{(k)}) - \rho(\lambda)(\alpha - k)| \leq C, \quad \forall k \leq \alpha - 1.$$

Consequently,

$$\sum_{k \leq \alpha-1} M(\lambda; ge^{-2\kappa k} \hat{q}^{(k)}) = \rho(\lambda) \sum_{k \leq \alpha-1} (\alpha - k) + O(\alpha) \\ = \frac{1}{2} \rho(\lambda) \alpha^2 + O(\alpha).$$

On the other hand, by (5.5)

$$M(\lambda; \pm gq_k) \leq Cg^{1/2}e^{-\kappa k} = Ce^{\kappa(\alpha-k)},$$

so that

$$(8.4) \quad \sum_{k > \alpha-1} M(\lambda; \pm gq_k) \leq C.$$

The asymptotics (5.10) follows.

The estimates (5.11) and (5.12) are proved in the same way. By (6.9) and (8.4), (4.11)

$$\widetilde{N}(\lambda_1, \lambda_2; \mathbf{A}_{gq}) \leq \sum_{k \leq \alpha-1} C + C' \leq C\alpha.$$

Furthermore, as $\hat{q}^{(k)}$ satisfies Condition 5.7, by (6.18) and (8.4), (4.11)

$$\begin{aligned}\tilde{N}(\lambda_1, \lambda_2; \mathbf{A}_{-gq}) &\leq \sum_{k \leq \alpha-1} N(\lambda_1, \lambda_2; -ge^{-2\kappa k} \hat{q}_k) + C' \\ &\leq \sum_{k \leq \alpha-1} C(\alpha - k) + C' \leq C'' \alpha^2.\end{aligned}$$

□

9. ASYMPTOTICS OF $M(\lambda, \mathbf{A}_{\pm gq})$ AND $N(\lambda, \mathbf{A}_{\pm gq})$

Here we turn to the study of the sums (4.1) and (4.2). As before, to ensure that they are finite we assume that $\lambda, \lambda_1, \lambda_2$ satisfy (4.21) with the same closed interval I . Due to the presence of exponential terms in the sums, their study is more complicated than that of $\tilde{M}$, $\tilde{N}$, and hence the asymptotic formulae are less explicit. Another feature is that for the exponential and power-like potentials the results are qualitatively different.

9.1. Exponential potentials. In this subsection we always (except for Theorem 9.3) assume that

$$(9.1) \quad q(t) = Q(t) = e^{-2\kappa t}.$$

This assumption allows one to obtain asymptotic formulae based on the “self-similarity” property of the function $e^{-2\kappa t}$. Introduce the notation

$$\ln b = \beta > 0.$$

Theorem 9.1. *Assume (9.1). Then the following two statements hold:*

- (i) *Let $\kappa > 0$ be arbitrary. Then there exist functions $\varphi_{\pm}$ that are 2κ -periodic, bounded and separated from zero, such that*

$$(9.2) \quad \lim \left[g^{-\frac{\beta}{2\kappa}} N(\lambda_1, \lambda_2; \mathbf{A}_{\pm gq}) - \varphi_{\pm}(\ln g) \right] = 0, \quad g \rightarrow \infty.$$

- (ii) *Suppose that $\kappa > 0$ is arbitrary if $\mathfrak{s} = +1$ and $\kappa < \beta$ if $\mathfrak{s} = -1$. Then there exist two functions $\psi_{\pm}$ that are 2κ -periodic, bounded and separated from zero, such that*

$$(9.3) \quad \lim \left[g^{-\frac{\beta}{2\kappa}} M(\lambda, \mathbf{A}_{\pm gq}) - \psi_{\pm}(\ln g) \right] = 0, \quad g \rightarrow \infty.$$

We precede the proof with an elementary but convenient lemma:

Lemma 9.2. *Let $n(t), t \in \mathbb{R}$ be a bounded function such that*

$$(9.4) \quad \begin{cases} n(t) = 0 & \text{for all } t \leq t_0 \text{ with some } t_0 > 0, \\ n(t) \leq Ct^{\frac{\beta}{2\kappa} - \epsilon}, t \geq t_0, & \text{for some } \epsilon > 0. \end{cases}$$

Then for the function

$$(9.5) \quad N(g) = n(g) + (1 - b^{-1}) \sum_{k \geq 1} e^{\beta k} n(ge^{-2\kappa k})$$

there exists a function ϕ which is 2κ -periodic, bounded and separated from zero, such that

$$\lim[g^{-\frac{\beta}{2\kappa}}N(g) - \phi(\ln g)] = 0, \quad g \rightarrow \infty.$$

Proof. The sum in the r.h.s. of (9.5) is finite, since for sufficiently large k we have $ge^{-2\kappa k} \leq t_0$. Denote

$$\xi(g) = g^{-\frac{\beta}{2\kappa}}n(g), \quad \Xi(g) = g^{-\frac{\beta}{2\kappa}}N(g).$$

Then (9.5) yields

$$\Xi(g) = \xi(g) + (1 - b^{-1}) \sum_{k \geq 1} \xi(ge^{-2\kappa k}),$$

which in its turn implies that

$$\Xi(g) - \Xi(ge^{-2\kappa}) = \xi(g) - b^{-1}\xi(ge^{-2\kappa}).$$

Using the notation (6.1) and introducing new functions $F(\alpha) = \Xi(g)$, $f(\alpha) = \xi(g)$, we arrive at the equation

$$F(\alpha) - F(\alpha - 1) = f(\alpha) - b^{-1}f(\alpha - 1).$$

Since $n(t)$ satisfies (9.4), $f(\alpha) = 0$ for $\alpha \leq \alpha_0 = (2\kappa)^{-1} \ln t_0$, and $f(\alpha) \leq Ce^{-2\kappa\epsilon\alpha}$, $\alpha \geq \alpha_0$. Therefore all the conditions of the Renewal Theorem are satisfied (see [14], Chapter XI.1, or a modern exposition in [16]), which guarantees the existence of a 1-periodic function $\tilde{\phi}$ which is bounded and separated from zero, such that

$$F(\alpha) = \tilde{\phi}(\alpha) + o(1), \quad \alpha \rightarrow \infty.$$

which leads to (9.2) after substitution $\phi(\alpha) = \tilde{\phi}(\alpha/(2\kappa))$. □

Proof of Theorem 9.1. (i) The proof is done simultaneously for both signs $\mathfrak{s} = \pm 1$. We use Lemma 9.2 with $n(g) = N(\lambda_1, \lambda_2; \pm gq)$. Since $N(\lambda_1, \lambda_2; \pm gq_k) = n(ge^{-2\kappa k})$, by (4.2) the function $N(g)$ in the r.h.s. of (9.5) coincides with $N(\lambda_1, \lambda_2; \mathbf{A}_{\pm gq})$. By (5.5) and (4.11) $n(t) = 0$, $t \leq t_0$ for a sufficiently small $t_0 > 0$. Moreover, by (6.18) or (6.7), the second condition in (9.4) is also fulfilled for any $\epsilon < \beta(2\kappa)^{-1}$. Thus the required asymptotics (9.2) follows from Lemma 9.2.

(ii) The cases $\mathfrak{s} = +1$ and $\mathfrak{s} = -1$ are treated separately.

Let first $\mathfrak{s} = +1$. Denote now $n(g) = M(\lambda; gq)$. Then, similarly to the first part of the proof, the total counting function (4.1) coincides with (9.5). By (5.5) and (6.6) the function n satisfies (9.4) for any $\epsilon < \beta(2\kappa)^{-1}$. Thus Lemma 9.2 guarantees the asymptotics (9.3) for $\mathfrak{s} = +1$.

In the case $\mathfrak{s} = -1$, $\kappa < \beta$, the first condition in (9.4) is satisfied for $n(g) = M(\lambda; -gq)$ in view of (5.5). Besides, (5.5) ensures also that the second condition is satisfied with $\epsilon = \beta(2\kappa)^{-1} - 1/2 > 0$. Again, Lemma 9.2 leads to (9.3) for $\mathfrak{s} = -1$. □

For $\mathfrak{s} = -1$ the cases $\varkappa < \beta$ and $\varkappa > \beta$ are described by Theorem 9.1 and Lemma 5.3 respectively. Let us handle the critical case $\beta = \varkappa$. We emphasise that this is the only asymptotic formula in this subsection which does not require the exact equality $q(t) = e^{-2\varkappa t}$.

Theorem 9.3. *Suppose that*

$$q(t) = Q(t)(1 + o(1)), \quad t \rightarrow \infty, \quad Q(t) = e^{-2\varkappa t},$$

with $\varkappa = \beta$, and that q satisfies Condition 5.7. Then

$$\lim_{g \rightarrow \infty} \frac{M(\lambda; \mathbf{A}_{-gq})}{g^{1/2} \ln g} = \frac{1 - b^{-1}}{2\pi \varkappa^2},$$

Proof. By (5.5) $M(\lambda; -gq_k) = 0$ for all $k \geq \alpha + A$ with a sufficiently large A , and

$$\sum_{k \leq A} b^k M(\lambda; -gq_k) + \sum_{\alpha - A \leq k \leq \alpha + A} b^k M(\lambda; -gq_k) \leq CA g^{1/2}.$$

Consequently

$$(9.6) \quad \lim_{g \rightarrow \infty} \frac{M(\lambda; \mathbf{A}_{-gq})}{g^{1/2} \ln g} = \lim_{g \rightarrow \infty} \frac{1 - b^{-1}}{g^{1/2} \ln g} \sum_{A < k < \alpha - A} e^{\varkappa k} M(\lambda; -gq_k),$$

if the limit in the r.h.s. exists. For $k \in (A, \alpha - A)$ apply Theorem 6.6 and the relation (4.5) to obtain the asymptotics

$$\begin{aligned} M(\lambda; -gq_k) &= \frac{g^{1/2}}{\pi} \int_0^\infty \sqrt{q(k+t)} dt + O(\alpha) \\ &= g^{1/2} e^{-\varkappa k} ((\pi \varkappa)^{-1} + o_A(1)) + O(\alpha), \end{aligned}$$

where $o_A(1) \rightarrow 0$ as $A \rightarrow \infty$ uniformly in k, g . Therefore the sum in the r.h.s. of (9.6) equals

$$\begin{aligned} &g^{1/2} \sum_{A < k < \alpha - A} \frac{1}{\pi \varkappa} + g^{1/2} \alpha o_A(1) + O(\alpha) \sum_{A < k < \alpha - A} e^{\varkappa k} \\ &= \frac{g^{1/2} \alpha}{\pi \varkappa} + g^{1/2} O(A) + g^{1/2} \alpha o_A(1) + e^{\varkappa \alpha} e^{-\varkappa A} O(\alpha). \end{aligned}$$

Since $\alpha = \ln g / (2\varkappa)$, now it follows from (9.6) that

$$\limsup_{g \rightarrow \infty} \left| \frac{M(\lambda; \mathbf{A}_{-gq})}{g^{1/2} \ln g} - \frac{1}{2\pi \varkappa^2} \right| = o_A(1) + O(e^{-\varkappa A}).$$

Since A is arbitrary, the required result follows. $\square$

9.2. Power-like potentials. For power-like potentials the asymptotic formulae that we obtain, are less informative since they are established for $\ln M$ and $\ln N$. For the sake of illustration we consider here only $M(\lambda; \mathbf{A}_{\pm gq})$. The corresponding asymptotics of $N(\lambda; \mathbf{A}_{\pm gq})$ can be easily derived using the same argument as well. For simplicity we assume that $q(t) = Q(t) = (1+t)^{-2\gamma}$. For more general power-like potentials the results follow by monotonicity of M with respect to the potential. Recall that I denotes the interval defined in (4.21). We also use the notation $d_{\pm} = |\lambda - \lambda_{\mp}|$, $\sigma_{\pm} = d_{\pm}^{-\frac{1}{2\gamma}}$ introduced in Lemma 7.1(iii).

Theorem 9.4. *Let I be a closed interval defined in (4.21), which is strictly inside the gap (λ_-, λ_+) . Let $q(t) = (1+t)^{-2\gamma}$, $\gamma > 0$. Then for any $\lambda \in I$*

$$\lim \alpha^{-1} \ln M(\lambda; \mathbf{A}_{\pm gq}) = \beta d_{\pm}^{-\frac{1}{2\gamma}}, \quad g \rightarrow \infty,$$

where $\beta = \ln b$.

Proof. Upper bound. A straightforward perturbation argument ensures that $M(\lambda; \pm gq_k) = 0$ if $gq_k(t) \leq d_{\pm}$, $\forall t > 0$, i.e. for

$$k \geq K_1 = K_1(g) = (d^{-1}g)^{\frac{1}{2\gamma}} = \alpha d_{\pm}^{-\frac{1}{2\gamma}}.$$

It follows from Lemmas 5.2 and 6.2 that $M(\lambda; \pm gq_k) \leq Cg^{\omega}$ with some $\omega = \omega(\gamma) > 0$. Substituting this bound in (4.1), gives

$$\begin{aligned} M(\lambda; \mathbf{A}_{\pm gq}) &= M(\lambda; \pm qq) + (1 - b^{-1}) \sum_{k \in \mathbb{N}} b^k M(\lambda; \pm gq_k) \\ &\leq Cg^{\omega} \sum_{k=0}^{K_1} b^k \leq C'g^{\omega} b^{K_1}, \end{aligned}$$

with a constant C' depending only on b, γ .

Lower bound. For the lower bound we drop all but one term from the sum (4.1): for any $\epsilon \in (0, 1)$ we have

$$M(\lambda; \mathbf{A}_{\pm gq}) \geq (1 - b^{-1})b^k M(\lambda; \pm gq_k), \quad k = [(1 - \epsilon)K_1].$$

By Lemmas 7.3 and 7.1(iii),

$$M(\lambda; \pm gq_k) \geq c\alpha F_{\pm}(\lambda; \beta_k) \geq c'\alpha (d_{\pm}^{-\frac{1}{2\gamma}} - \beta_k)^{\frac{3}{2}},$$

for sufficiently large α . Since $\beta_k = (k+1)\alpha^{-1}$, we see that the r.h.s. is bounded from below by $c''\alpha\epsilon^{3/2}$ with a constant depending only on $d_{\pm}$. Consequently,

$$M(\lambda; \mathbf{A}_{\pm gq}) \geq c\epsilon^{\frac{3}{2}}\alpha b^{K_1(1-\epsilon)},$$

with a constant depending on b, λ and γ . Since $\epsilon > 0$ is arbitrary, in combination with the upper bound, this gives the required asymptotics. $\square$

10. ACKNOWLEDGEMENTS

The second author (M.S.) was partly supported by the Minerva center for non-linear physics and by the Israel Science Foundation, and partly by the EPSRC grant GR/N 37193/01.

This paper was essentially completed when M.S. was visiting King's College, London in April-May 2001. The authors are grateful to Yu. Safarov for valuable discussions.

REFERENCES

1. S. Alama, P.A. Deift, and R. Hempel, *Eigenvalue branches of the Schrödinger operator $H - \lambda W$ in a gap of $\sigma(H)$* , Comm. Math. Phys. **121** (1989), 291–321.
2. M.Sh. Birman, *Discrete spectrum in the gaps of a continuous one for perturbations with large coupling constant*, Estimates and asymptotics for discrete spectra of integral and differential equations (Leningrad, 1989–90), 57–73. Adv. Soviet Math., **7**, Amer. Math. Soc., Providence, RI, 1991.
3. M.Sh. Birman and V.V. Borzov, *On the asymptotics of the discrete spectrum for certain singular differential operators*, Probl. Mat. Fiz. **5** (1971), 24–38; English transl. in Topics in Math. Phys. **5** (1972).
4. M.Sh. Birman and A. Laptev, *Discrete spectrum of the perturbed Dirac operator*, Ark. Mat. **32** (1994), no. 1, 13–32.
5. ———, *The negative discrete spectrum of a two-dimensional Schrödinger operator*, Comm. Pure Appl. Math. **49** (1996), no. 9, 967–997.
6. M.Sh. Birman, A. Laptev, and M. Solomyak, *On the eigenvalue behaviour for a class of differential operators on semiaxis*, Math. Nachr. **195** (1998), 17–46.
7. M. Sh. Birman and M. Solomyak, *Quantitative analysis in Sobolev imbedding theorems and applications to spectral theory*, Tenth Mathem. School, Izd. Inst. Mat. Akad. Nauk Ukrain 5–189, SSSR, Kiev, 1974 (Russian); English transl. in Amer. Math. Soc. Translations, (2), **114** (1980), 1–132.
8. ———, *Estimates for the number of negative eigenvalues of the Schrödinger operator and its generalizations*, Estimates and asymptotics for discrete spectra of integral and differential equations (Leningrad, 1989–90), 1–55, Adv. Soviet Math., **7**, Amer. Math. Soc., Providence, RI, 1991.
9. ———, *On the negative discrete spectrum of a periodic elliptic operator in a waveguide-type domain, perturbed by a decaying potential*, J. d'Anal. Math. **83** (2000), 337–391.
10. M.Sh. Birman and G.D. Raikov, *Discrete spectrum in the gaps for perturbations of the magnetic Schrödinger operator*, Estimates and asymptotics for discrete spectra of integral and differential equations (Leningrad, 1989–90), 75–84, Adv. Soviet Math., **7**, Amer. Math. Soc., Providence, RI, 1991.
11. R. Carlson, *Hill's equation for a homogeneous tree*, Electron. J. Differen. Equations 1997 (23), 30 pp. (electronic).
12. ———, *Nonclassical Sturm–Liouville Problems and Schrödinger operators on radial trees*, Electron. J. Differen. Equations 2000 (71), 24 pp. (electronic).
13. P. A. Deift and R. Hempel, *On the existence of eigenvalues of the Schrödinger operator $H - \lambda W$ in a gap of $\sigma(H)$* , Comm. Math. Phys. **103** (1986), 461–490.
14. W. Feller, *An introduction to the probability theory and its applications*, Vol. II, John Wiley and Sons, Inc., New York-London-Sidney-Toronto, 1971.
15. M. Klaus, *On the point spectrum of Dirac operators*, Helv. Phys. Acta **53** (1980), 453–462.
16. M. Levitin and D. Vassiliev, *Spectral asymptotics, renewal theorem, and the Berry conjecture for a class of fractals*, Proc. London Math. Soc. (3) **72** (1996) 188–214.
17. K. Naimark and M. Solomyak, *Eigenvalue estimates for the weighted Laplacian on metric trees*, Proc. London Math. Soc. (3) **80** (2000) 690–724.

18. ———, *Geometry of the Sobolev spaces on the regular trees and Hardy's inequalities*, Russian Journal of Mathematical Physics, **8**, No 3 (2001).
19. A.V. Sobolev, *Weyl asymptotics for the discrete spectrum of the perturbed Hill operator*, Estimates and asymptotics for discrete spectra of integral and differential equations (Leningrad, 1989–90), 159–178, Adv. Soviet Math., **7**, Amer. Math. Soc., Providence, RI, 1991.

CENTRE FOR MATHEMATICAL ANALYSIS AND ITS APPLICATIONS, UNIVERSITY OF SUSSEX, FALMER,
BRIGHTON, BN1 9QH, UK

E-mail address: `A.V.Sobolev@sussex.ac.uk`

DEPARTMENT OF MATHEMATICS, WEIZMANN INSTITUTE, REHOVOT, ISRAEL

E-mail address: `solom@wisdom.weizmann.ac.il`